

PHYTOSOCIOLOGICAL STUDIES ON THE STEPPE COMMUNITIES OF NATURAL PASTURES IN ŞANLIURFA, UPPER MESOPOTAMIA, TURKEY

CEVHERI, C.^{1*} – ASLAN, M.² – ATAMOV, V.³

¹*Department of Biology, Faculty of Science and Art, Harran University, Şanlıurfa, Turkey*

²*Faculty of Education, Harran University, Şanlıurfa, Turkey*

³*Department of Biology, Faculty of Science and Art, Recep Tayyip Erdoğan University, Rize, Turkey*

**Corresponding author*

e-mail: ccevheri@harran.edu.tr; phone: +90-414-318-3566

(Received 28th May 2019; accepted 28th Nov 2019)

Abstract. This phytosociological study was carried out in the steppe of Fatik Mountains in Şanlıurfa, between the years 2002-2005. At the end of this study, it was underlined that the dominant vegetation type was steppe in the region and different steppe associations were determined. These were also; *Festuco callieri–Teucrietum poliae* ass. nova, *Astragalo strictifolii–Salvietum sclarea* ass. nova, *Phlomido bruguieri–Thymbretum spicatae* ass. nova, *Asphodelo aestivum–Teucrietum poliae* ass. nova, *Cynodo dactyloni–Bromoetum danthoniae* ass. nova, and *Prosopo farctae–Avenetum erianthae* ass. nova. During the floristic surveys, 153 taxa belonging to 26 families and 107 genera were identified. *Poaceae* with 31 taxa (20.3%), *Asteraceae* with 24 taxa (15.7%), *Fabaceae* with 22 (14.4%), and *Apiaceae* with 13 taxa (8.5%) were the richest families.

Keywords: *steppe vegetation, grazing, ecosystems, characteristic species, association*

Introduction

Turkey is located at the intersection of two important gene centers namely the Mediterranean and Near East and has the ninth highest biodiversity among terrestrial countries. Natural species of most cereals cultivated for human consumption grow on steppe ecosystems (Ture and Bocuk, 2007).

Fatik Mountain, located at the southwestern part of Şanlıurfa, was selected as the research area since it has a peculiar steppe vegetation and has been paid less attention so far as “Adıgüzel and Aytac, 2001” mentioned earlier. The studied area is in the C7 square according to grid system adopted by Davis (1988). Phytogeographically located in the Irano–Turanian Floristic Region, extending between the longitudes of 37.2 North and 37.3 and between the latitudes of 38.6 and 38.7 East (Anonymous, 1995).

The central, eastern and southeastern parts of Turkey belong phytogeographically to the Irano–Turanian region (Zohary, 1973). The Irano–Turanian region is separated into two parts the West and the East Irano–Turanian subregions. There are three provinces in Turkey belonging to the West Irano–Turanian subregion namely; Central Anatolian, Irano–Anatolian and the Mesopotamian provinces. According to this classification, Southeast Anatolia is located in the Mesopotamian province of the West Irano–Turanian subregion (Zohary, 1973).

According to Flora of Davis (Davis 1965-1985; Davis et al.; 1988; Güner et al., 2000) some researchers among them Kotschy, Sintenis, Davis, and Zohary have visited and

gathered plant specimens from Şanlıurfa by Baytop (2003). Some floristic and taxonomic studies were carried out in and around Southeast Anatolia (Kaynak and Ketenoğlu, 1980; Malyer, 1981-1983; Kaynak, 1987-1989a, b; Yıldırım, 1992-1994; Yıldız and Aktoklu, 1996; Ertekin and Saya, 1997; Ertekin, 2002; Adıgüzel and Aytac, 2001; Aslan and Turkmen, 2001-2003; Turkmen et al., 2002; Akan et al., 2005; Aydoğdu and Akan, 2005). But there has been no vegetational research data except for a study carried out by Zohary (1973). In addition, Southeast Anatolia is one of the most little known or unknown regions of Turkey (Davis, 1965-1985; Çırpıcı, 1987; Donner, 1990).

In the region, the natural areas are used as pastures like in all arid countries. Artificial pasture culture has not been demanded yet. It has been stated that it contains wheat, beans in pastures in arid regions and includes drought-resistant different vegetation types and other herbaceous plants such as Savanne, Praerie, Steppe, Maguis, Phyrwana. (Gençkan, 1983). This understanding and using of pastures are applied in Şanlıurfa on large scale, like in the most of regions of Turkey. In this region the open places which are out of using, belong to state, like slopes, stony, rocky places, stream beds, hills etc. (Çullu et al., 2002). Şanlıurfa is one of little researched region in Turkey (Aslan, 2015). It has 724.529 ha of area. 263.507 ha of this area is used as natural pastures. In the region, the pastures have the poorest plant cover (10-15%) in Turkey. 0.82 ha of area is divided for each cow. In the past years there have been significant changes in the diversity of plants in this area due to random overgrazing and in some years fires (steppe fires) due to the abundance of animals. But, when the pastures of the region are taken into consideration it is necessary to divide 4 ha area for each cow. This shows that the pastures are used five times excessively. Excessive and irregular using have caused succession and degradation. The primary vegetation has transformed to secondary vegetation and converting towards semi-desert ecosystem (Atamov et al., 2004).

The characteristic vegetation type is steppe in this region (Atamov et al., 2004). The threat factors are excessive grazing and transforming of natural pastures to agricultural areas. Because of increased industrial activities, urbanization, tourism activities, creating new agricultural areas, mining activities, using agricultural methods and overgrazing, the natural structures of steppe ecosystems have been getting destroyed (Ture and Bocuk, 2007). 74 taxa are under threat and of these 10 taxa on Karacadağ, 13 taxa in Ceylanpınar are endemic (Özhatay et al., 2003; Aslan, 2018). In Şanlıurfa, the steppe vegetation is dominant because of ecological, geographical and antropogenic factors. Usually, animals eat legumes, grasses and similar crops, which are fresh and delicious for a period of one year. However, the perennial shrub, smelly and prickly plants are not preferred.

Under effects of topographic and edafic factors, different steppe vegetation types are found in the region. In this steppe, wheats are dominant and different plants can be found, which have life forms like geophytes, camephytes, cryptophytes, hemicryptophytes and terophytes. In some districts, some remains of forest and rare old trees are found. In addition, the efemerics which grow rapidly and complete their life cycles in short time after spring and autumn rains are present in the vegetation.

The constant plants are perennial herbs, shrubs and trees that are seen rarely (Atamov et al., 2007). But, the efemerics are found and affect quality of fodder plants and yields positively.

In the region, watery agriculture has been made for years. Agricultural method applied to irrigation in semi-arid and arid regions depending on the characteristics of the plant sown or to obtain higher yields. Since these areas are not natural areas, weed yield

is considerably reduced due to the excess number of animals. Holophytic areas have risen 394%. The agriculture of wheat and cotton have been stopped in some places because of salinity and these areas have been used as pastures for the last 15 years (Çullu et al., 2002).

Materials and methods

The city Şanlıurfa in Southeast of Anatolia is between 37° 49' 12" – 40° 10' 00" East meridian and 36° 41' 28" – 37° 55' 50" North parallel. In Şanlıurfa the uneven areas are at percentage of 60.4%, mountains 22%, plains 16.3% and plateaus 1.3%. Harran plain, covering the study area, is one of the most important agricultural areas of Turkey and located between 36°47' and 39°15' east longitude and 36°40' and 37°41' north latitude within the borders of the province of Şanlıurfa in Southeastern Anatolian region. The most important plains are named Harran, Suruç, Viranşehir, stayed on south part of Şanlıurfa. Ceylanpınar plain are stayed on southwest. In addition, Halfeti, Hilvan and Bozova plains have large areas. The average altitude is 375 m. The lowest plain is Harran plain. Viranşehir plain is at east, Suruç plain is at west of Harran plain. It is 141 535 ha (Anonymous, 1995) (Figs. 1, 2 and 3).

Floristic and phytosociologic research studies were carried out between the years 2000-2005 at Karacadağ, Tektek, Fatik, Kaşmer, Nemrut, Kalecik mountains and Direkli Hills where are used as natural pastures. Regular geobotanical trips were organized. Plant specimens were collected and preserved according to the herbarium techniques. These were identified with the help of "Flora of Turkey and the East Aegean Islands" (Davis, 1965-1985, 1988; Güner et al., 2000) and other relevant publications (Akman et al., 2001) as well as the flora of Syria, Iraq and Iran. In addition, the plants of Birecik Dam lake area and halophytic plants in Akçakale were gathered and classified (Ertekin and Saya, 1997; Ertekin, 2002; Adıgüzel and Aytac, 2001; Aslan, 2002; Aslan and Türkmen, 2001-2003; Kaya, 2002; Atamov et al., 2004; Akan et al., 2005; Aydın, 2003; Aydoğdu and Akan, 2005; Ayalp, 2005; Atamov et al., 2005).




Figure 1. Location of the study area in Turkey




Figure 2. The map of Fatik Mountains



Figure 3. A photo of the research area

Braun-Blanquet's (1964) method was used for the naming of phytosociological plant groups. Plants were sampled twice a month during certain vegetation periods and plant samples were taken to identify the plants and examine the necessary materials.) These plants have been protected in the Herbarium of Harran University. The climatic data were taken from the Şanlıurfa Meteorology Station and has been evaluated according to these data. Approximately 500 g. soil samples were taken from 5 cm under the soil in the research areas in spring, summer and autumn. The soil samples were dried naturally and analysis of them were made by the Ministry of Agriculture and the Rural Affairs General Directorate of Rural Services Şanlıurfa Research Institute.

Results and discussion

As it is seen in *Figure 4* and *Table 1*, the climate in Şanlıurfa is arid from June to October in a long period. In this period, most annual plants die. Only perennial herbs, shrubs and trees are alive. Semiarid Mediterranean climate occurs in research area. According to Emberger the precipitation–temperature coefficient (Q) is 42.94 (Akman, 1990). Annual mean temperature is 18.7 °C. The maximum mean temperature (M) is 46.8 °C, in July. The minimum mean temperature (m) is -6.8 °C, in February. Annual rainfall is about 457.8 mm (Anonymous, 2001) and the seasonal precipitation regime is winter, spring, autumn and summer. This is the first variant of the East Mediterranean precipitation regime. The ombrothermic diagram shows dry and rainy periods (*Fig. 4*).

Table 1. The bioclimate and fall regime of Sanliurfa

Altitude (m)	P (mm)	M (°C)	m (°C)	Q	PE	S	Fall regime	Bioclimate
547	457.8	46.8	-6.8	42.94	7.2	0.18	W.Sp.A.Su	Semi-arid, cold in winter




Figure 4. The climatic diagram of Şanlıurfa. **a:** Meteorology station; **b:** The elevation of the Meteorology station (m); **c:** The year of temperature and precipitation; **d:** The mean annual temperature (°C); **e:** The mean annual precipitation (mm); **f:** The curved line of temperature; **g:** The curved line of precipitation; **h:** Arid season; **i:** Humid season; **m:** The least mean temperature of the most cold month (°C); **n:** Absolute minimum temperature (°C); **r:** Probable frosty month

In the different plant associations in the steppe vegetation, the average productivity changes between 1.0-7.0 kg m⁻². Excessive aridity and temperature have caused decreasing productivity and quality of fodder plants in pastures. From the end of autumn to the middle of spring, the ephemerals grow and these plants increase the quality and productivity of fodders by 2-3 times. At the end, yield of natural pastures, number of fodder plant species and their populations have been reduced day to day because of excessive and unconscious grazing, migration, negative habitat conditions like aridity and temperature. The plants which have prickles, etheric oils, latex, poison, bitter and fluff have spread out. Excessive grazing has caused the transforming of the primary steppe to secondary and then semi-desert to desert vegetation.

In most of the places in this area, the covering degree of the flora is 40-50%. And this is the characteristic property of half-desert and desert type associations. The flora

has generally made up of a single layer and the average height of the grass covering is 10-20 cm.

On the contrary, the number of plants which are not eaten by animals have been increased. Thus, the primary steppe vegetation has been transformed to secondary and it is seen that the succession has been transformed to semi-desert conditions.

Immigration is widespread in this region and emigrants migrate from East Anatolia to use pastures for grazing. Thousands of sheep destroy the all immature and senile groups of plants. The yield of pastures has been decreased. This situation has caused transforming from steppe to desertation.

In the flora of Şanlıurfa, 71 families, 798 taxa belonging to 342 genera have been determined (Atamov et al., 2004). 135 of these plants were monocotyledonae, 663 were dicotyledonae.

The number of plants belonging to *Poaceae* family were more than others in spite of having less species (56) than *Fabaceae* (101). The dominant taxa which spread out in naturel pastures were as follows: *Bromus japonicus* Thunb. subsp. *japonicus*, *B. squarrosus* L., *B. sterilis* L., *B. tectorum* L., *Avena sterilis* L. subsp. *sterilis*, *Aegilops triuncialis* L. subsp. *triuncialis*, *A. neglecta* Req. ex Bertol, *Festuca callieri* (Hackel ex St. – Yves) F. Markgraf subsp. *callieri*, *Hordeum murinum* L. subsp. *glaucum* (Steudel) Tzvelev.

Some plants belonging to *Fabaceae*, have fodder quality. Most of them are used as fodders. These were as follows; *Astragalus onobrychis* L., *A. aleppicus* Boiss., *A. immarginatus* Labm., *A. dactylocarpus* Boiss., *Medicago rigidula* (L.) var. *rigidula*, *Onobrychis caput-galli* (L.) Lam. *O. crista-galli* (L.) Lam. *O. galegifolia* Boiss. *O. kotschyana* Fenzl. *Trifolium scabrum* L., *T. speciosum* Willd., *T. dasyurum* C. Presl, *T. angustifolium* L. var. *intermedium*, *T. boissieri* Guss. ex Boiss., *T. meironense* Zoh., *T. campestre* Schreb., *Trigonella monspeliaca* L., *T. mesopotamica* Hub. Mor., *Vicia mollis* Boiss., *Coronilla orientalis* Miller var. *orientalis*, *C. scorpioides* (L.) Koch, *Hedysarum varium* Willd.

In the research area, 153 taxa belonging to 26 families and 107 genera were identified. *Poaceae* with 31 taxa (20.3%), *Asteraceae* with 24 taxa (15.7%), *Fabaceae* with 22 (14.4%), and *Apiaceae* with 13 taxa (8.5%) were the richest families.

Braun-Blanquet's (1964) method was used for the naming of phytosociological plant groups.

32 sample parcels were evaluated and 6 new associations and their upper categories are as follows:

Astragalo–Bromotea Quezel 1973

Astragalo–Bromotalia Quezel 1973

Agropyro–Stachyon Quezel 1973

1. *Festuco callieri–Teucrietum poliae* ass. ova

Onobrychido armenae–Thymetalia leucostomi Akman, Ketenoglu Quezel 1984

2. *Astragalo strictifolii–Salvietum sclarea* ass. nova

3. *Phlomido bruguieri–Thymbretum spicatae* ass. nova

4. *Asphodelo aestivum–Teucrietum poliae* ass. nova

5. *Cynodo dactyloni–Bromoetum danthoniae* ass. nova

6. *Prosopo farctae–Avenetum erianthae* ass. nova

***Festuco callieri–Teucrietum poliae* ass. nova**

There were no trees and shrubs in its structure. Only grass layer was present and covering was 40-80%. The length was between 10 and 35 cm (Table 2).

Festuco callieri–Teucrietum poliae association was spreading out where the calcereous main rock, gravelled and small pit areas. According to chemical and physical properties of soils where the association was spreading out homogenous, pH was between 7.0 and 7.71, the rate of lime (CaCO₃) was 3.8%, phosphorus (P₂O₅) was 5.2-5.3 kg dec⁻¹; potassium (K₂O) was 118.8-308.8 kg dec⁻¹ and organic component was 2.17-2.48% (Table 3).

Table 2. *Festuco callieri–Teucrietum poliumae* ass. nova. Thypus: Sample Parcel 62

Sample parcel No	12	23	34	35	37	41	42	46	48	50	Frequency
Square size (m ²)	50	50	50	50	50	50	50	50	50	50	
Altitude (m)	500	478	530	540	570	510	515	530	535	550	
Gradient (%)	20	30	30	35	30	25	30	35	30	25	
Direction	N	NW	NW	N	N	N	N	N	N	NW	
Length of cover (cm)	10	25	20	17	20	22	20	25	25	20	
Number of species	21	21	20	16	17	18	18	15	16	17	
Cover (%)	60	55	55	60	65	65	60	60	60	65	
Characteristic species of Association											
<i>Festuca callieri</i>	22	22	+2	+2	+1	22	22	33	22	11	V
<i>Teucrium polium</i>	–	22	+1	11	11	1	33	33	+1	11	IV
<i>Trifolium tomentosum</i>	+2	+2	+1	+2	+2	III
Characteristic species of Allians <i>Festuca–Teucrium</i>											
<i>Phleum phleoides</i>	+1	+2	+2	+2	.	.	.	+2	+2	+2	III
<i>Phlomis bruguieri</i>	+2	+2	+2	+2	+2	+2	+2	+1	.	.	III
<i>Eryngium creticum</i>	+1	+2	+2	.	.	+2	+1	.	.	.	II
Characteristic species of Ordo <i>Daphno–Festucetalia</i>											
<i>Thymbra spicata</i> var. <i>spicata</i>	22	22	22	11	11	+2	+2	+2	.	.	III
<i>Lolium rigidum</i>	+2	+2	.	.	.	+2	+1	+1	+1	+1	II
<i>Hordeum murinum</i> subsp. <i>glaucum</i>	+2	.	.	+2	+2	+2	+2	+2	+2	.	III
Characteristic species of Classis <i>Daphno–Festucetea</i>											
<i>Salvia sclarea</i>	.	.	.	+2	+2	+2	+2	+2	+2	+1	II
<i>Teucrium polium</i>	+2	+1	+2	+2	+2	II
<i>Scutellaria tomentosa</i>	+2	+1	+2	+2	+2	II
<i>Aegilops cylindrica</i>	+2	+2	+1	+1	+1	+1	+1	.	.	.	II
Participants											
<i>Allium flavum</i> subsp. <i>tauricum</i> var. <i>tauricum</i>	+2	+2	+2	+2	+2	+2	+2	+2	+2	+2	III
<i>Avena sterilis</i> subsp. <i>sterilis</i>	+1	+1	.	.	.	+2	+2	+2	+2	+1	II
<i>Euphorbia denticulata</i>	+1	+1	+1	+1	+1	+1	I
<i>Tordylium syriacum</i>	+1	+2	+2	+2	+1	I
<i>Erodium cicutarium</i> subsp. <i>cutarium</i>	+1	+1	+1	I
<i>Cichorium glandulosum</i>	+1	+1	+2	+2	.	.	I
<i>Crepis reuterana</i> subsp. <i>reuterana</i>	+1	+1	+1	I
<i>Gagea luteoides</i>	.	+1	I

<i>Thlaspi perfoliatum</i>	+1	+1	+1	+1	I
<i>Muscari comosum</i>	+1	+1	+1	.	I
<i>Scandix pecten-veneris</i>	+1	+1	.	.	+1	I
<i>Lappula barbata</i>	+1	I
<i>Cynosurus echinatus</i>	.	+1	I
<i>Torilis tenella</i>	+1	+1	+1	.	I
<i>Hypocoum imberbe</i>	.	.	+1	+1	I
<i>Senecio vulgaris</i>	+1	I
<i>Anthemis haussknechtii</i>	+1	+1	+1	.	.	+1	I
<i>Gypsophila antari</i>	+1	+2	+2	+2	.	.	.	+1	.	.	II
<i>Medicago minima</i> var. <i>minima</i>	+1	+1	+2	+2	+2	.	.	.	+2	+2	II
<i>Echinops microcephalus</i>	.	.	+1	I
<i>Centaurea solstitialis</i> subsp. <i>solstitialis</i>	+1	I
<i>Hypocoum pendulum</i>	+1	+1	I
<i>Torilis leptophylla</i>	.	.	+2	+1	+1	+1	+1	.	.	.	I

Table 3. Chemical characteristics of the soil of the research area

Location	Depth (cm)	Saturation with water (%)	Total salt (%)	pH	CaCO ₃ (%)	P ₂ O ₅	K ₂ O	Organic mass (%)
Ugurlu Village	0-20	77	0.071	7.70	3.8	5.3	118.8	2.48
Kocoren Village	0-20	77	0.060	7.71	3.8	5.2	308.8	2.17
Ikizce Village	0-20	68	0.073	7.28	2.6	4.9	179.2	1.44
Altun Village	0-20	66	0.098	7.42	6.8	4.5	108	2.35

The characteristic species of the association were *Festuca callieri* subsp. *callieri*, *Teucrium polium*. Each species are the elements of Irano–Turanian phytogeographic region.

Classis: *Daphno–Festucetea* Quezel 1964

Ordo: *Daphno–Festucetalia* Quezel 1964

Alliance: *Festuca–Teucrium* ass. nova

Asphodelo aestivi–Teucrietum poliae ass. nova

The characteristic species of the association were *Asphodelus aestivus* and *Teucrium polium*. This association was at the north part of the Fatik Mountains and at the environs of Stone quarry. 10 sample areas were used. The gradient was 35-40%, The length was between 600 and 700 m (Table 4).

Table 4. *Asphodelo aestivi–Teucrietum poliae* ass. nova. *Thypus*: Sample Parsel 63

Sample parcel No	60	61	62	67	70	73	77	79	81	82	Frequency
Square size (m ²)	50	50	50	50	50	50	50	50	50	50	
Altitude (m)	531	535	500	510	515	520	530	505	500	517	
Gradient (%)	30	20	25	25	30	30	35	30	35	35	
Direction	W	W	W	W	NW	NW	NW	NW	NW	W	

Length of cover (cm)	20	25	20	20	25	15	20	25	20	30	
Number of species	14	13	11	8	12	13	13	10	11	11	
Cover (%)	55	50	45	55	50	55	60	50	40	45	
Characteristic species of Association											
<i>Astragalus strictifolius</i>	11	+2	+2	11	11	12	22	22	22	+2	V
<i>Teucrium polium</i>	+2	22	22	22	11	.	.	11	+2	+2	VI
<i>Asphodelus aestivus</i>	12	33	33	.	.	+2	+2	.	.	.	III
Characteristic species of Allians <i>Agropyro–Stachyon</i>											
<i>Euphorbia denticulata</i>	+1	.	+1	.	+1	.	+2	+2	+2	.	III
<i>Centaurea solstitialis</i> subsp. <i>solstitialis</i>	.	+2	.	.	+2	+2	.	.	.	+1	II
<i>Bromus japonicus</i> subsp. <i>japonicus</i>	+1	+2	.	.	33	+1	+2	+1	.	.	III
Characteristic species of Ordo <i>Astragalo–Bromotalia</i>											
<i>Astragalus lamarckii</i>	11	+2	+2	22	22	11	11	11	+2	+2	IV
<i>Bromus danthoniae</i>	+2	+2	+1	+1	.	.	.	+1	+1	+2	III
<i>Scutellaria tomentosa</i>	+1	+1	+2	+1	+1	+1	II
Characteristic species of Classis <i>Astragalo–Bromotea</i>											
<i>Arrhenatherum kotschyi</i>	+2	+2	.	.	+2	+2	.	.	+1	11	II
<i>Phlomis kurdica</i>	+2	.	.	+2	+1	+1	.	+1	.	.	II
<i>Gypsophila antari</i>	+1	.	+1	.	.	.	+1	.	.	.	I
<i>Stipa holosericea</i>	.	.	.	+1	+2	+2	+2	+2	.	.	II
Participants											
<i>Poa bulbosa</i>	+1	+1	+2	+2	+	+1	+2	+2	+2	+2	II
<i>Trifolium speciosum</i>	.	.	+2	.	+2	+2	+1	.	.	+2	II
<i>Torilis arvensis</i>	+1	+1	+1	+1	+2	+1	II
<i>Anthemis haussknechtii</i>	+1	+1	+1	.	.	.	+1	.	.	.	I
<i>Lolium rigidum</i>	.	.	.	+1	+1	+2	+2	.	+2	.	I
<i>Capparis spinosa</i>	+1	+1	.	.	.	I
<i>Convolvulus arvensis</i>	+1	+2	I

The association contain one layer composed of grass and average length was between 20 and 30 cm. The plant cover was changing between 40 and 60%.

Classis: *Astragalo–Bromotea* Quezel 1973

Ordo: *Astragalo–Bromotalia* Quezel 1973

Alliance: *Agropyro–Stachyon* Quezel 1973

This association was spreading out on the calcereous main rock, and gravelled and small pit areas. According to chemical and physical properties of soils, pH was: 7.28-7.71, the rate of lime (CaCO₃) was 3.8%, phosphorus (P₂O₅) was 5.2 kg/dec.; potassium (K₂O) was 118.8 kg/dec. and organic component was 2.48%. (Table 3). The yield of pasture was low, between 0.5 and 1.0 ha⁻¹, because of excessive grazing. 20 taxa were determined.

Cynodo dactyloni–Brometum danthoniae ass. nova

The characteristic species of the association were *Cynodon dactylon* and *Bromus danthoni*. It was defined in stream beds and depressed places on the southwest of İközce village at Fatik mountains. The gradient was low (5-10%), and the altitude was between

600 and 700 m. 10 sample parcels were used. This association composed of only grass layer and the average length was between 10 and 20 cm., covering was 60–65% and taxon number was poor (between 7 and 13) (Table 5).

Classis: *Astragalo–Bromotea* Quezel 1973

Ordo: *Astragalo–Bromotalia* Quezel 1973

Alliance: *Agropyro–Stachyon* Quezel 1973

This association was spread out at red-brown soils. According to chemical and physical properties of soils; pH was: 7.42, the rate of lime (CaCO₃) was 6.8%, phosphorus (P₂O₅) was 4.5 kg dec⁻¹; potassium (K₂O) was 108.0 kg dec⁻¹. and organic component was found as 2.35% (Table 3). The yield of pasture was 1.0 ha⁻¹

The association *Cynodo dactyloni–Brometum danthoniae* is composed of 27 taxa (Table 5).

Table 5. *Cynodo dactyloni–Bromoetum danthoniae* ass. nova. *Thypus*: Sample Parcel 64

Sample parcel No	**	63	64	69	71	74	75	76	78	80	Frequency
Square size (m ²)	50	50	50	50	50	50	50	50	50	50	
Altitude (m)	470	485	480	485	490	485	480	458	500	505	
Gradient (%)	10	20	25	15	20	25	15	20	25	20	
Direction	NE	NE	NE	NE	NE	N	N	N	NE	NE	
Length of cover (cm)	15	15	10	10	15	15	10	15	10	20	
Number of species	13	13	13	12	13	8	12	12	7	10	
Cover (%)	65	60	65	65	60	60	65	55	55	65	
Characteristic species of Association											
<i>Cynodon dactylon</i>	.	+2	22	33	33	22	23	23	+2	+2	V
<i>Bromus danthoniae</i>	+1	+2	+1	+2	+2	11	11	22	11	12	IV
<i>Bromus japonicus</i>	+2	+1	+2	11	11	11	+2	.	.	.	III
Characteristic species of Allians <i>Agropyro–Stachyon</i>											
<i>Avena eriantha</i>	+1	+1	+1	+2	+2	+2	+2	.	.	.	III
<i>Eryngium creticum</i>	+1	.	+1	.	+1	.	.	+1	+1	+1	II
<i>Cichorium glandulosum</i>	.	.	.	+1	.	.	.	+1	.	+1	I
Characteristic species of Ordo <i>Astragalo–Bromotalia</i>											
<i>Astragalus lamarckii</i>	.	.	+2	+1	+1	+2	+2	+2	+2	+2	III
<i>Bromus rubens</i>	+2	+2	+2	.	.	.	+1	+1	.	.	II
<i>Euphorbia characias</i> subsp. <i>wulfenii</i>	.	.	.	+1	+1	+1	I
Characteristic species of Classis <i>Astragalo–Bromotea</i>											
<i>Aegilops cylindrica</i>	+2	+2	+1	+2	.	II
<i>Hordeum bulbosum</i>	.	.	+1	+2	+2	+2	I
<i>Ziziphora capitata</i>	+1	+1	+1	+2	I
<i>Scrophularia canina</i> subsp. <i>bicolor</i>	+1	+1	+1	.	.	.	I
Participants											
<i>Scandix pecten–veneris</i>	.	.	.	+1	+1	+1	+2	+2	+2	+2	III
<i>Verbascum kotschyi</i>	.	+1	+1	+1	I
<i>Centaurea consanguinea</i>	.	.	+1	.	.	.	+1	+1	+1	.	I
<i>Veronica beccabunga</i> subsp. <i>beccabunga</i>	+1	+1	.	.	+1	I
<i>Ranunculus arvensis</i>	11	+1	+2	+2	+2	+1	II
<i>Secale sylvestris</i>	+1	+1	+1	.	I

<i>Allium flavum</i> subsp. <i>tauricum</i> var. <i>tauricum</i>	+1	+1	+1	I
<i>Senecio vulgaris</i>	+1	+1	.	.	.	I
<i>Crepis reuterana</i> subsp. <i>reuterana</i>	+1	.	.	.	+1	+1	I
<i>Trifolium tomentosum</i>	+1	.	I
<i>Hypocoum pendulum</i>	.	.	.	+1	+2	+2	I
<i>Avena sterilis</i> subsp. <i>sterilis</i>	+1	+1	+1	I
<i>Hypericum capitatum</i> var. <i>capitatum</i>	+1	I

Phlomido bruguieri–Thymbretum spicatae ass. nova

Phlomis bruguieri and *Thymbra spicata* subsp. *Spicata* were the characteristic species of this association. It was spread out on the south part of the Fatik mountain, between İkizce village and Altun mezra village. The gradient was 15-30% and the altitude was between 610 and 850 m (Table 6).

The association contained only one layer like the others, grass layer. The average length was 12-25 cm and rate of covering was 50-60% (Table 6).

Classis: *Astragalo–Bromotea* Quezel 1973

Ordo: *Onobrychido armenae–Thymetalia leucostomi* Akman, Ketenoğlu Quezel 1984

Alliance: *Phlomido armeniaceae–Astragalion microcephali* Akman, Ketenoğlu Quezel 1986

It was spreading out at red-brown soils. According to chemical and physical properties of soils, pH was: 7.70, the rate of lime (CaCO₃) was 3.8%, phosphorus (P₂O₅) was 5.3 kg dec⁻¹; potassium (K₂O) was 118.8 kg dec⁻¹ and organic component was found as 2.48% (Table 3). The yield of pasture was 1.0-1.5 ha⁻¹ 29 taxa were determined from this association (Table 6).

Table 6. *Phlomido bruguieri–Thymbretum spicatae* ass. nova *Thypus*: Sample Parcel 17

Sample parcel No	10	13	14	15	17	19	21	22	28	30	33	65	66	Frequency	
Square size (m ²)	50	50	50	50	50	50	50	50	50	50	50	50	50		
Altitude (m)	610	605	707	710	715	720	725	810	815	750	820	822	815		
Gradient (%)	20	15	25	20	25	35	30	20	20	25	30	30	30		
Direction	SE	S	S	S	S	SE	SE	SE	SE	SE	SE	SE	S		
Length of cover (cm)	25	20	20	25	20	20	25	20	25	20	20	20	25		
Number of species	12	12	10	11	15	9	9	9	12	7	9	9	7		
Cover (%)	50	55	55	60	60	60	65	55	55	50	50	60	55		
Characteristic species of Association															
<i>Phlomis bruguieri</i>	+2	+2	22	22	11	11	22	22	11	+2	+2	+2	+1	V	
<i>Thymbra spicata</i>	+2	+2	+2	+1	+1	11	11	11	+2	+2	+1	+1	+2	IV	
Characteristic species of Allians <i>Phlomido armeniaceae–Astragalion microcephali</i>															
<i>Hordeum murinum</i> subsp. <i>glaucum</i>	+2	+2	22	22	11	11	11	+1	+2	+2	22	11	11	IV	
<i>Phleum phleoides</i>	+1	+2	+2	+2	+2	+2	.	.	III	
<i>Teucrium parviflorum</i>	+2	.	.	.	+2	.	.	.	+2	II	
<i>Euphorbia characias</i> subsp. <i>wulfenii</i>	+2	+2	.	.	+2	.	.	I	

Characteristic species of Ordo <i>Onobrychido armenae–Thymetalia leucostomi</i>														
<i>Onobrychis caput-galli</i>	+2	+2	+1	+2	+2	+2	+2	+2	+2	.	.	+2	+2	IV
<i>Bromus scoparius</i>	.	+2	+2	+2	+1	+1	+1	+1	+1	+2	+2	+1	.	III
<i>Medicago rigidula</i>	+1	+1	+1	+1	.	.	.	I
Characteristic species of Classis <i>Astragalo–Bromotea</i>														
<i>Bromus japonicus</i>	.	+2	+2	+2	+1	+1	+1	+2	+2	+2	+2	+2	+2	III
<i>Satureja hortensis</i>	+1	+1	+1	+2	+2	+2	+2	II
<i>Scabiosa calocephala</i>	+1	+1	+1	+2	+2	+2	.	.	II
Participants														
<i>Hypocoum imberbe</i>	+1	I
<i>Torilis tenella</i>	+1	I
<i>Xanthium strumarium</i> subsp. <i>strumarium</i>	+1	.	.	.	+1	.	I
<i>Cynosurus echinatus</i>	+1	+1	I
<i>Eremopyrum bonaepartis</i> subsp. <i>bonaepartis</i>	.	+1	.	.	+1	I
<i>Urginea maritima</i>	+1	.	.	I
<i>Secale sylvestris</i>	+1	.	I
<i>Gagea luteoides</i>	+2	I
<i>Trifolium stellatum</i>	+1	I
<i>Crepis reuterana</i> subsp. <i>reuterana</i>	+1	I
<i>Sedum caespitosum</i>	.	.	+1	+1	I
<i>Echinops microcephalus</i>	+1	.	.	+1	+1	I
<i>Eryngium creticum</i>	+1	.	.	+1	+1	I
<i>Capsella bursa-pastoris</i>	+1	.	.	.	+1	.	.	.	I
<i>Thlaspi perfoliatum</i>	+1	+1	.	+1	I
<i>Velezia rigida</i>	+1	.	.	I

Prosopo farctae–Avenetum erianthae ass. nova

The characteristic species were *Prosopis farcta*, *Avena eriantha* and *Bromus scoparius*. It was determined from the Northwest of Küçük Güneş, Aşağı Güneş and Yukarı Güneş villages of Fatik Mountain at 450-500 m altitudes.

The association composed of only grass layer except *Prosopis farcta*, the average length was between 15 and 30 cm and the covering was between 50 and 65%. The number of taxon was between 12 and 18 (Table 7).

Classis: *Astragalo–Bromotea* Quezel 1973

Ordo: *Astragalo–Bromotalia* Quezel 1973

Alliance: *Prosopo–Bromuon* ass. nova

This association was spreading out at red-brown soils. According to chemical and physical properties of soils, pH was: 7.28, the rate of lime (CaCO₃) was 2.6%, phosphorus (P₂O₅) was 4.9 kg dec⁻¹; potassium (K₂O) was 179.2 kg dec⁻¹ and organic component was measured as 1.44% (Table 3). The yield of pasture was 1.0-1.5 ha⁻¹ 26 taxa were determined (Table 7).

Table 7. *Prosopio farctae–Avenetum erianthae* ass. nova *Thypus*: Sample Parcel 9

Sample parcel No	2	5	6	8	9	11	12	16	18	20	25	26	27	Frequency	
Square size (m ²)	50	50	50	50	50	50	50	50	50	50	50	50	50		
Altitude (m)	450	455	460	470	475	480	485	490	495	500	505	510	515		
Gradient (%)	10	15	10	10	10	10	15	15	15	10	10	10	15		
Direction	E	E	E	SE	SE	SE	SE	SE	SE	SE	SE	SE	SE		
Length of cover (cm)	25	20	20	25	30	15	15	25	15	15	25	25	20		
Number of species	18	13	12	13	13	12	12	13	13	12	13	13	14		
Cover (%)	55	50	55	50	50	45	50	60	55	50	45	50	50		
Characteristic species of Association															
<i>Prosopis farcta</i>	+2	+2	33	33	22	22	11	11	11	22	22	33	33	V	
<i>Bromus scoparius</i>	+1	+1	+2	.	.	22	22	+2	+2	+1	+1	+2	22	IV	
<i>Avena eriantha</i>	+2	+2	22	22	22	+2	+2	+2	+2	+2	+2	.	+1	IV	
Characteristic species of Allians <i>Prosopo–Bromuon</i> ***															
<i>Phlomis bruguieri</i>	+2	+2	+2	+2	+2	+1	+1	+2	+2	+2	+2	+2	+2	III	
<i>Hordeum murinum</i> subsp. <i>glaucum</i>	+1	+1	+2	+2	11	11	11	11	11	+1	+1	.	.	III	
<i>Salvia sclarea</i>	+1	.	.	.	+1	+1	+1	+2	+2	+2	+2	.	.	II	
Characteristic species of Ordo <i>Astragalo–Bromotalia</i>															
<i>Astragalus lamarckii</i>	+1	+2	+2	+2	+2	+2	+2	+2	.	.	+2	.	.	II	
<i>Bromus japonicus</i>	+1	+1	11	+2	+1	.	.	.	+1	+2	.	+1	12	II	
<i>Juncus articulatum</i>	+1	+1	+1	22	22	22	22	.	.	.	11	11	.	III	
Characteristic species of Classis <i>Astragalo–Bromotea</i>															
<i>Phleum phleoides</i>	+2	+2	+2	+2	+1	+1	+1	+2	+2	+1	+1	+1	+1	II	
<i>Phlomis bruguieri</i>	+2	+2	.	.	+2	+2	.	.	+2	+2	+2	.	.	II	
<i>Cichorium glandulosum</i>	+1	.	+1	+1	.	.	.	+1	+1	+1	.	.	.	I	
Participants															
<i>Althea officinalis</i>	+1	+1	+1	+1	+1	.	+2	+2	+2	II	
<i>Verbascum kotschyi</i>	.	.	.	+1	+2	+2	+1	+1	+1	.	.	+1	+1	II	
<i>Linum pubescens</i> subsp. <i>pubescens</i>	+1	.	.	+1	+1	.	.	I	
<i>Euphorbia szovitsii</i> var. <i>szovitsii</i>	+1	+1	+1	.	.	I	
<i>Eryngium campestre</i>	.	+1	+1	+1	I	
<i>Hypericum capitatum</i> var. <i>capitatum</i>	+1	+1	.	.	.	+1	.	I	
<i>Trifolium tomentosum</i>	+1	+1	.	I	
<i>Hordeum bulbosum</i>	+1	+1	I	
<i>Aegilops cylindrica</i>	.	.	.	+1	.	+1	+1	.	I	
<i>Torilis arvensis</i>	+1	.	.	.	I	
<i>Carduus pycnocephalus</i> subsp. <i>albidus</i>	+1	+1	.	+1	I	
<i>Anthemis haussknechtii</i>	.	+1	+1	+1	I	
<i>Erodium cicutarium</i> subsp. <i>cutarium</i>	+1	+1	+1	I	
<i>Lappula barbata</i>	.	.	+1	+1	.	.	.	+1	+1	I	
<i>Secale sylvestre</i>	.	.	.	+1	+1	I	

Astragalo strictifolii–Salvietum sclarea ass. nova

Astragalus strictifolius var. *strictifolius* and *Salvia sclarea* were the characteristic taxa of this association. It was determined between Fatik and İkizce villages at northeast direction. The altitude was 600-750 m and the gradient was between 20 and 35%. The association composed of only grass layer except *Astragalus strictifolius* var. *strictifolius*. The average length was between 20 and 25 cm and the covering was between 50 and 65%. The number of taxon was between 10 and 18 (Table 8).

Classis: *Astragalo–Bromote* Quezel 1973

Ordo: *Astragalo–Bromotalia* Quezel 1973

Alliance: *Agropyro–Stachyon* Quezel 1973

This association was spreading out at the red-brown soils on the arid stone. According to chemical and physical properties of soils, pH was: 7.42, the rate of lime (CaCO₃) was 6.8%, phosphorus (P₂O₅) was 4.5 kg dec⁻¹; potassium (K₂O) was 108.0 kg dec⁻¹ and organic component was measured as 2.35% (Table 3). The yield of dry grass was between 300 and 400 g m⁻² 35 taxa were determined (Table 8).

Table 8. *Astragalo strictifolii–Salvietum sclarea ass. nova* Thypus: Sample Parcel 36

Sample parcel No	1	3	4	36	38	44	45	47	49	51	52	54	58	Frequency	
Altitude (m)	50	50	50	50	50	50	50	50	50	50	50	50	50		
Gradient (%)	510	520	530	540	550	555	557	560	565	567	570	571	572		
Direction	30	35	20	20	25	30	30	35	35	30	35	30	25		
Length of cover (cm)	SE	SE	SE	S	S	S	SW	SW	SW	S	S	S	S		
Number of species	25	20	20	25	30	35	35	30	30	25	30	25	25		
Cover (%)	13	12	15	16	16	13	13	9	12	10	15	12	12		
Altitude (m)	50	55	55	60	65	60	60	55	55	60	60	60	50		
Characteristic species of Association															
<i>Astragalus strictifolius</i> var. <i>strictifolius</i>	12	12	22	+1	+1	22	+1	+2	22	22	22	11	11	IV	
<i>Salvia sclarea</i>	12	.	12	11	+1	+1	+2	+2	11	22	+2	+2	11	IV	
Characteristic species of <i>Allians Agropyro Stachyon</i>															
<i>Thymra spicata</i>	.	.	+2	22	33	+2	+2	+2	+2	22	22	11	11	IV	
<i>Phlomis kurdica</i>	+2	+2	+2	+2	+2	+2	+2	.	III	
<i>Euphorbia characias</i> subsp. <i>wullfenii</i>	.	.	+2	+2	+2	+2	+2	+2	II	
<i>Stipa holosericea</i>	+2	+2	+2	.	.	II	
Characteristic species of Ordo <i>Astragalo–Bromotalia</i>															
<i>Festuca callieri</i> subsp. <i>callieri</i>	+2	+2	+2	+2	+2	+2	II	
<i>Astragalus lamarckii</i>	22	+2	+2	22	22	.	.	.	22	11	11	22	11	IV	
<i>Phlomis</i> ssp.	+2	+2	+2	.	.	.	+2	+2	+2	.	+2	.	.	III	
Characteristic species of Classis <i>Astragalo–Bromotea</i>															
<i>Eryngium creticum</i>	.	.	+2	+2	+2	+2	+2	.	.	+2	+2	.	+2	III	
<i>Teucrium polium</i>	+2	+2	.	.	+2	+2	+2	+2	II	
<i>Trifolium speciosum</i>	.	.	.	+2	+2	+2	+2	+2	+2	II	
<i>Ranunculus arvensis</i>	+2	.	+2	.	.	.	+1	+1	+1	+1	.	.	.	I	

	Participants													
<i>Eminium rauwolffii</i> var. <i>rauwolffii</i>	+2	+2	+2	+2	+2	+2	+2	+1	+1	+1	.	.	+2	III
<i>Muscari comosum</i>	+2	+2	+2	+1	+1	+1	.	II
<i>Gagea luteoides</i>	+2	+2	+2	+1	+2	+2	+1	+1	III
<i>Erodium cicutarium</i> subsp. <i>cutarium</i>	+1	+2	.	.	+1	+1	+1	.	.	I
<i>Achillea vermicularis</i>	+2	I
<i>Eryngium campestre</i>	.	.	.	+1	+2	+2	+1	I
<i>Capsella bursa-pastoris</i>	+1	+1	+1	I
<i>Avena sterilis</i> subsp. <i>sterilis</i>	+2	+1	+1	+2	+1	.	.	.	+2	I
<i>Echinaria capitata</i>	+1	+1	+1	+1	I
<i>Althea officinalis</i>	.	.	.	+2	.	.	+1	+2	I
<i>Veronica beccabunga</i> subsp. <i>beccabunga</i>	.	+1	+1	+1	+1	I
<i>Vulpia persica</i>	+1	+1	I
<i>Bromus japonicus</i>	.	.	.	+1	+1	+1	+1	.	.	I
<i>Eremopyrum bonaepartis</i> subsp. <i>bonaepartis</i>	+2	.	.	.	+2	.	.	I
<i>Convolvulus arvensis</i>	.	.	+2	+1	I
<i>Cichorium glandulosum</i>	+1	I
<i>Linum pubescens</i>	.	.	.	+1	.	.	.	+1	+1	I
<i>Aegilops cylindrica</i>	+2	.	.	.	+2	I
<i>Xanthium strumarium</i> subsp. <i>strumarium</i>	.	+1	+1	.	.	+1	.	.	I

Conclusion

In the region, due to excessive and uncontrolled grazing, steppe vegetation has turned into secondary vegetation rather than semi-desert and desert-type vegetation. Therefore, these areas should be controlled and over-grazing should be prevented. In natural pastures, the value of the forage plant and the most suitable pasture capacity should be determined. The productivity of pastures is related to the composition of plant species and habitat factors (Braun-Blanquet, 1964). Morphological structures and ecological habitat forms are one of the most important subjects that should be studied in pasture research projects.

First, the quality and quantity of a pasture should be examined. In the region, plant sociological characteristics of the dominant steppe vegetation change during the seasons. The efemers change the productivity and quality of the pastures. In pastures, forage crops have been reduced due to overgrazing, others spread if inedible to animals. These areas have been abandoned for agriculture and have become barren due to overgrazing. In the future research of these areas, controlled pasture stocking should be carried out together with controlled irrigation for the formation of productive pastures, and even pasture forage crops suitable for barren areas should be selected and trials should be conducted in the area. National and regional pasture management plans should continue to be implemented effectively, sustainable productivity must be ensured and these areas should be put back into service of Turkish livestock.

REFERENCES

- [1] Adıgüzel, N., Aytaç, Z. (2001): Flora of Ceylanpınar State Farm (Şanlıurfa-Turkey). – *Fl Medit* 11: 333-361.
- [2] Akan, H., Kaya, Ö. F., Eker, I., Cevheri, C. (2005): The Flora of Kaşmer Mountain, Sanliurfa, Turkey. – *Turk. J. Bot.* 29: 1-20.
- [3] Akman, Y. (1990): Climate and Bioclimate. – Palme Publications, Ankara.
- [4] Akman, Y., Ketenoglu, O., Geven, F. (2001): Vegetation Ecology and Research Methods. – Ankara University Publications No: 9, Ankara.
- [5] Anonymous (1995): Şanlıurfa Province Land Asset. – T. C. General Directorate of Rural Services Directorate Publications, Ankara.
- [6] Anonymous (2001): Meteorological Bulletin. – T. C. State Meteorological Service General Directorate No: A-09: 27-29, Ankara.
- [7] Aslan, M. (2002): Birecik Dam Lake area plants, hazard classes and conservation. – PhD Thesis, Cukurova University, Institute of Science and Technology, Adana.
- [8] Aslan, M. (2015): Succession of steppe areas after fire in the Gap Region of Turkey. – *Bangladesh J. Bot.* 44(4): 489-497.
- [9] Aslan, M. (2018): Habitats change in Ceylanpınar State's farm and the dangerous classes of plants (Şanlıurfa-Turkey). – *International Journal of Science and Research (IJSR)* 7(11): 1766-1772.
- [10] Aslan, M., Türkmen, N. (2001): New floristic records for C7 grid square. – *Herb Systematic Botanic Journal* 8(2): 69-73.
- [11] Aslan, M., Türkmen, N. (2003): New floristic records for squares C6 and C7 from Turkey. – *Herb Systematic Botanic Journal* 10(2): 163-168.
- [12] Atamov, V., Cevheri, C., Parmaksız, A. (2004): Şanlıurfa Natural Rangeland Forage Bitkileri. – National Biology Congress XVII: 120, Adana, Turkey.
- [13] Atamov, V., Yavuz, M., Cevheri, C., Parmaksız, A., Çen, O., Çetin, E. (2005): "Halophytes and Halophidication In Harran Plain (ŞANLIURFA)". – *Biosalina Agriculture & High Salinity Tolerance*, 9-14 January 2005, Muğla, Turkey.
- [14] Atamov, V., Aslan, M., Cevheri, C., Cetin, E. (2007): Contribution to the flora of Fatik Mountain. – *Asian J. Plant Sci.* 6(1): 1-11.
- [15] Ayalp, G. (2005): Flora and vegetation of Mezra Town. – Master Thesis, Harran University Institute of Science and Technology, Şanlıurfa.
- [16] Aydın, N. (2003): Masted Hills flora. – Master Thesis. Harran University Institute of Science and Technology, Şanlıurfa.
- [17] Aydoğdu, M., Akan, H. (2005): The flora of Kalecik Mountain, Şanlıurfa-Turkey. – *Turk. J. Bot.* 29: 155-174.
- [18] Baytop, A. (2003): History of Botanical Studies in Turkey. – Çetin Printing, İstanbul.
- [19] Braun-Blanquet, J. (1964): *Pflanzensoziologie*. Dritte Aufl. – Springer Verlag, Wien.
- [20] Çırpıcı, A. (1987): The studies on the flora and vegetation of Turkey. – *Turk. J. Bot.* 11(2): 217-232.
- [21] Çullu, M., Almaca, A., Sahin, Y., Aydemir, S. (2002): Application of GIS monitoring soil salinization in The Harran Plain. – *International Conference on Sustainable Land Use and Management*, Şanlıurfa, Turkey, pp. 326-333.
- [22] Davis, P. H. (ed). (1965-1985): *Flora of Turkey and the East Aegean Islands*. Vol. 1-9. – Edinburgh University Press, Edinburgh.
- [23] Davis, P. H. (1988): *Flora of Turkey and the East Aegean Islands*. Vol. 10. – Edinburgh University Press, Edinburgh.
- [24] Donner, J. (1990): Distribution Maps to P. H. Davis "Flora of Turkey, 1-10". – *Linzer Biol. Beitr.* 22(2): 381-515.
- [25] Ertekin, S. (2002): *Karacadağ Plant Diversity Book*. – Sustainable Rural and Urban Development Association Publication, Diyarbakır.

- [26] Ertekin, S., Saya, Ö. (1997): New floristic records for the various grid squares from the Fabaceae. – Tr J of Botany 21: 187-189.
- [27] Gençkan, M. (1983): Agriculture of Forage Crops. – Ege University Printing House, Bornova, Izmir.
- [28] Güner, A., Özhatay, N., Ekim, T., Baser, K. H. C. (2000): Flora of Turkey and East Aegean Islands. Vol. 11. – Edinburgh University Press, Edinburgh.
- [29] Kaya, F. Ö (2002): Flora of Tektik Mountains (Sanliurfa). – Master Thesis, Harran University Institute of Science and Technology, Sanliurfa.
- [30] Kaynak, G. (1987): New record for some squares in flora of Turkey. – Nature TU Bot. Magazine 11(2): 118-123.
- [31] Kaynak, G. (1989a): Contribution to the flora of Karacadağ (Urfa and Diyarbakır Provinces). – Nature TU Bot. Magazine 13(3): 376-397.
- [32] Kaynak, G. (1989b): Ecological and chorological investigations on the ferns of Diyarbakır and surrounding provinces. – Nature TU Bot. Magazine 13(3): 437-451.
- [33] Kaynak, G., Ketenoğlu, O. (1980): New fern specimens in South East Anatolian Region. – Ist Univ Science Faculty Mec Series B 45: 199-202.
- [34] Malyer, H. (1981): A chorological study on the geophytes of the Iridaceae family of the Diyarbakır region. – Nature Science Journal Basic B 6: 17-20.
- [35] Malyer, H. (1983): Chorological and ecological investigations on geophytes of Liliaceae and Iridaceae families in Karacadağ (between Diyarbakır and Urfa). – Nature Science Journal Basic, Series C 7(3): 279-288.
- [36] Özhatay, N., Byfield, A., Atay, S. (2003): Turkey's Important Plant Alanları. – WWF-Turkey Natural Life Protection Foundation, Istanbul.
- [37] Ture, C., Bocuk, H. (2007): An investigation on the diversity, distribution and conservation of Poaceae species growing naturally in Eskisehir province (Central Anatolia-Turkey). – Pak. J. Bot 39(4): 1055-1070.
- [38] Türkmen, N., Aslan, M., Düzenli, A. (2002): New records for the various squares in the flora of Turkey. – Journal of Herb Systematic Botany 9(2): 63-68.
- [39] Yıldırım, Ş. (1992): Distribution of new plant to Turkey from various squares. – Nature TU Bot Derg 16: 207-214.
- [40] Yıldırım, Ş. (1994): From Turkey Brassicaceae (Cruciferae) were from New Floris squares for the various family Kayıtlar. – T J of Botany 18: 389-392.
- [41] Yıldız, B., Aktoklu, E. (1996): New floristic records from C7 square (Malatya-Adıyaman). – Tr J of Botany 20: 207-211.
- [42] Zohary, M. (1973): Geobotanical Foundations in the Middle East. Vol. 1-2. – Gustav Fischer Verlag, Stuttgart.