



Cloning, expression and biochemical characterization of a β -carbonic anhydrase from the soil bacterium *Enterobacter* sp. B13

Ayşenur Eminoğlu, Daniela Vullo, Aycan Aşık, Dilşat Nigar Çolak, Claudiu T. Supuran, Sabriye Çanakçı & Ali Osman Beldüz

To cite this article: Ayşenur Eminoğlu, Daniela Vullo, Aycan Aşık, Dilşat Nigar Çolak, Claudiu T. Supuran, Sabriye Çanakçı & Ali Osman Beldüz (2016) Cloning, expression and biochemical characterization of a β -carbonic anhydrase from the soil bacterium *Enterobacter* sp. B13, Journal of Enzyme Inhibition and Medicinal Chemistry, 31:6, 1111-1118, DOI: [10.3109/14756366.2015.1100176](https://doi.org/10.3109/14756366.2015.1100176)

To link to this article: <https://doi.org/10.3109/14756366.2015.1100176>



Published online: 26 Oct 2015.



Submit your article to this journal [↗](#)



Article views: 1211



View related articles [↗](#)



View Crossmark data [↗](#)



Citing articles: 4 View citing articles [↗](#)

RESEARCH ARTICLE

Cloning, expression and biochemical characterization of a β -carbonic anhydrase from the soil bacterium *Enterobacter* sp. B13

Ayşenur Eminoğlu¹, Daniela Vullo², Aycan Aşık³, Dilşat Nigar Çolak⁴, Claudiu T. Supuran², Sabriye Çanakçı⁵, and Ali Osman Beldüz⁵

¹Department of Biology, Faculty of Art and Science, Molecular Biology Research Laboratories, Recep Tayyip Erdogan University, Rize, Turkey, ²Neurofarba Department, Sezione Di Scienze Farmaceutiche E Nutraceutiche, University of Florence, Sesto Fiorentino (Firenze), Italy, ³Department of Medical Biology, Faculty of Medicine, Selcuk University, Konya, Turkey, ⁴Bulancak School of Applied Sciences, Giresun University, Giresun, Turkey, and ⁵Department of Biology, Faculty of Science, Karadeniz Technical University, Trabzon, Turkey

Abstract

A recombinant carbonic anhydrase (CA, EC 4.2.1.1) from the soil-dwelling bacterium *Enterobacter* sp. B13 was cloned and purified by Co²⁺ affinity chromatography. Bioinformatic analysis showed that the new enzyme (denominated here B13-CA) belongs to the β -class CAs and to possess 95% homology with the ortholog enzyme from *Escherichia coli* encoded by the *can* gene, whereas its sequence homology with the other such enzyme from *E. coli* (encoded by the *cynT* gene) was of 33%. B13-CA was characterized kinetically as a catalyst for carbon dioxide hydration to bicarbonate and protons. The enzyme shows a significant catalytic activity, with the following kinetic parameters at 20°C and pH of 8.3: k_{cat} of $4.8 \times 10^5 \text{ s}^{-1}$ and k_{cat}/K_m of $5.6 \times 10^7 \text{ M}^{-1} \times \text{s}^{-1}$. This activity was potently inhibited by acetazolamide which showed a K_i of 78.9 nM. Although only this compound was investigated for the moment as B13-CA inhibitor, further studies may reveal new classes of inhibitors/activators of this enzyme which may show biomedical or environmental applications, considering the possible role of this enzyme in CaCO₃ biomineralization processes.

Keywords

β -Carbonic anhydrase, *Enterobacter* sp., kinetics

History

Received 26 August 2015
Revised 21 September 2015
Accepted 22 September 2015
Published online 22 October 2015

Introduction

Carbonic anhydrases (CAs) [EC 4.2.1.1] are Zn(II)-dependent metalloenzymes which catalyze the reversible hydration of CO₂ to HCO₃⁻ and H⁺. CAs have important functions in the eukaryotes as well as in prokaryotes, such as pH regulation, biosynthetic reactions, respiration, CO₂ transport, photosynthesis, among others², and are known to be among the fastest enzymes described so far. Bacterial CAs have an essential role in the life cycle of these organisms, as well as in the regulation of calcium carbonate mineralization, at least in some bacterial species. Indeed, CaCO₃ mineralization by bacteria is crucial for soil and ground water remediation, sequestration and capture of atmospheric CO₂, as well as sand and soil strengthening and consolidation³.

CAs are encoded by six evolutionary unrelated families, the α , β , γ , δ , ζ and η -CAs⁵, which are present in organisms all over the phylogenetic tree. Among them, the α -, β -, γ - and δ -CAs are broadly distributed in many microorganisms¹, whereas the η -CAs were described only in protozoa for the moment⁵.

To date, two different β -CAs have been reported in *Escherichia coli*, the most investigated bacterium for the molecular biology viewpoint. One such enzymes is encoded

by the *cynT* gene and was the first β -CA identified in this bacterium, as a product of the *cyn* operon, being shown to possess sequence homology with higher plants β -CAs and to function in the process of cyanate utilization as a nitrogen source by this bacterium^{6,7}. The second one is *can* (previously called *yadF*), which has 30% amino acid sequence homology with *CynT*⁸. It was reported that *can* is essential for the growth of *E. coli* under atmospheric CO₂ conditions². It has been also shown that expression of *can* is susceptible to the increase of cell density and to the rise in the medium temperature. In *E. coli*, CA activity is normally provided by the expression of both these enzymes, although detailed data regarding their localization within the bacterium are missing for the moment. In fact, bicarbonate/CO₂ are necessary during the normal growth of this bacterium and for the biosynthesis of several biomolecules involved in vital functions. It has been suggested that *can* is conserved throughout *Enterobacteriaceae* genera (the type of bacteria to which *E. coli* also belongs)⁹. However, only four β -CA genes have been identified in *Enterobacteriaceae* so far, i.e. *cynT*, *can* (previously *yadF*), *cah* and *mig-5*⁹. Recently, β -CAs from two *Enterobacter* species were phylogenetically compared with the *E. coli* CAs¹⁰, but only one of these enzymes has been investigated in detail for the moment, i.e. the one encoded by the *cynT* gene, for which the X-ray crystal structure was reported by Cronk et al.⁸

Although ultimately there are many studies on various bacterial β -CAs, the investigation of *Enterobacteriaceae* is

Address for correspondence: Claudiu T. Supuran, Neurofarba Department, University of Florence, Via Ugo Schiff 6, Polo Scientifico, Sesto Fiorentino (Firenze) 50019, Italy. Tel: +39-055-4573005. Fax: +39-055-4573729. E-mail: claudiu.supuran@unifi.it




Figure 1. Phylogenetic analysis of 16S rDNA similarities of *Enterobacter* sp. B13 based on the BLAST result using the neighbor-joining method. Scale bar represents 0.01 substitutions per nucleotide position. The organisms and GeneBank accession numbers of analyzed sequences are given in parenthesis.

rather limited with only one *E. coli* enzyme being investigated in some detail, as mentioned above. Here, we describe a new β -CA from a bacterium belonging to this bacteria family, more precisely from an *Enterobacter* sp. B13 recently isolated from soil in Northern Turkey.

Materials and methods

Sampling site, culture conditions and microorganisms

The bacterial specimens were isolated from soil and mud samples which were collected from a riverside nearby Trabzon, Sürmene, Turkey. The samples were diluted with dH₂O and filtrated. The filtrate inoculated in TSA plates and incubated at different temperatures. Selected colonies which grew at 37 °C were then inoculated to modified B-4 agar plates to monitor the CaCO₃ decomposition properties of the grown bacteria. Among these isolates, clone B13 showed CO₂ hydratase activity¹¹ and was subsequently used as a source of genomic DNA in this study.

Identification of the bacterial strain by 16S rRNA analysis

For identification of the strain, the 16S rRNA gene was amplified via PCR from genomic DNA with eubacterial universal primers and subsequently sequenced (Macrogen Inc., Seoul, Korea).

Cloning of *Enterobacter* sp. B13 *can* and construction of the recombinant expression vector

The genomic DNA of *Enterobacter* sp. was obtained with Wizard Genomic DNA Purification Kit (Promega, Madison, WI) and was used as a template for the CA gene. The amplification process was performed by PCR with two primers in order to amplify the *Enterobacter* sp. B13 *can* with *Nde*I (Biolab) and *Bam*HI (Thermo Scientific, Waltham, MA) recognition sites (*CANde*I; 5'-CCA TAT GAA CGA CAT AGA TAC-3' and *Bam*HI CAR5'-CGG ATC CTT ATT TAT GGT TNA CGT GC-3') with the following conditions: initial denaturation steps at 95 °C for 3 min, denaturation at 95 °C for 45 s, followed by annealing at 51 °C for 1 min and primer extension at 72 °C for 1 min 20 s, followed by a step at 72 °C for 5 min, for 35 cycles. The PCR products were cloned into pGEM-T Easy vector (Promega) and sequenced (Macrogen Inc., Seoul, Korea). After the sequence verification, both the positive clone and the expression vector pET-15b (Novagen) were digested with *Nde*I-*Bam*HI, then excised from the agarose gel (Qiaquick Gel Extraction Kit, Qiagen, Venlo, The Netherlands) and ligated (Thermo Scientific T4 Ligase, Thermo Scientific, Waltham, MA).

Construction and cloning of the recombinant plasmid was carried out in *E. coli* JM101 (Novagen, Podenzano, Italy) as described in the literature¹². Transformants were checked by enzymatic digestion.




Figure 2. SDS-PAGE analysis (15%) of *Enterobacter* sp. B13-CA (stained with coomassie brilliant blue). Lane 1, protein molecular weight markers (NEB, P7710S), lane 2, purified *Enterobacter* sp. B13-CA.

Overexpression in *E. coli* BL21 (DE3) *pLysS* and purification of the β -CA

The recombinant vector designed as *pB13 can* was transformed into *E. coli* BL21 (DE3)*pLysS* (Novagen) for overexpression and sequenced for correctness. A single colony, harboring the recombinant plasmid was chosen and cultured overnight in Luria Broth (LB) medium containing 50 μ g/ml ampicillin. This culture was used to inoculate 200 ml LB-ampicillin medium and incubated at 37 °C with vigorous shaking. When OD_{600} reached the mid-log phase, ~ 0.6 – 0.8 , 0.5 mM $ZnSO_4$ were added to the culture and expression was induced by 0.1 mM isopropyl β -D-thiogalactopyranoside (IPTG) for 24 h at 22 °C, for the production of recombinant B13-CA. Cells were then collected by centrifugation at 9000 rpm for 10 min at 4 °C. The pellet was resuspended in the lysis buffer (30 mM Tris- SO_4 (pH 8.0), 0.1% Triton X-100, 200 mM NaCl, 20 mM imidazole and 0.2 mg/ml lysozyme) and sonicated. After centrifugation at 7000 rpm for 20 min at 4 °C, the supernatant containing *N*-terminally His-tagged protein was purified by Co^{2+} (TALON Metal Affinity Resin, Clontech), pre-equilibrated with the wash buffer (50 mM NaH_2PO_4 (pH 8.0), 0.1% Triton X-100, 300 mM NaCl and 20 mM imidazole). After transferring, the lysate was incubated in the His beads in 15 ml falcon tubes overnight at +4 °C. Following the two washing

steps at pH 8.0 and one at pH 7.0 by gentle shaking at +4 °C, the CA was eluted with 250 mM imidazole in the same buffer (pH 7.0). SDS-PAGE (15%) was run to evaluate the purity of the protein. Purified B13-CA was dialyzed overnight in dialysis buffer (30 mM Tris- SO_4 , pH 8.0, 0.5 mM $ZnSO_4$, 1 mM dithiothreitol-DTT) at +4 °C. After dialysis, the concentration of the protein was measured with both NanoDrop Spectrophotometer 2000 and at 595 nm, with bovine serum albumin (BSA) as a standard.

Zymography

Zymography analysis of the CA activity was performed as described in the literature¹³. To determine the esterase activity, zymogram staining was performed with minor modifications as described previously¹⁴ at two different temperatures (+4 °C and also at RT) on SDS-PAGE (15%). In brief, followed by electrophoresis the gels were incubated in 100 mM Tris-HCl (pH 7.5) including 0.5% Triton X-100, for 6 h at +4 °C and for 4 h at RT. After incubation, the gels were rinsed in 100 mM Tris-HCl, containing 100 mM α -naphthyl acetate (α -NAc), dissolved in 2 ml acetone, and 100 mM Tris-HCl (pH 7.5) buffer containing 20 mg Fast Red Salt (two solutions were added to the gels at the same time) until the bands become visible.

Bioinformatic analysis

The nucleotide sequence of *Enterobacter* sp. B13 *can* was submitted to GeneBank under accession number KT184504. The identification of 16S rDNA sequence was determined by the EzTaxon identification tool (<http://www.ezbiocloud.net/eztaxon>) against 16S rRNA sequences found in the database, i.e. strains with validly published prokaryotic names¹⁵. *Haemophilus influenzae* ATCC was selected as outgroup member. Amino acid resemblance was determined by using the DELTA-BLAST server (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>).

For predicting the estimated molecular weight and to perform the multiple amino acid sequence alignment, Compute pI/Mw tool of ExPASy¹⁶ and ClustalW¹⁷ were used, respectively. Protein family and conserved domains were identified by Conserved Domain Search Database (<http://www.ncbi.nlm.nih.gov/cdd>). Determination of evolutionary familiarities was performed by Mega 6¹⁸.

CO_2 hydrase activity assay of B13-CA

An Applied Photophysics stopped-flow instrument was used for assaying the CA catalyzed CO_2 hydration activity¹⁹. Phenol red (at 0.2 mM) was used as indicator, working at the absorbance maximum of 557 nm with 10 mM TRIS (pH 8.3) as buffer and 0.1 M $NaClO_4$ (for maintaining constant ionic strength), at 20 °C, following the CA-catalyzed CO_2 hydration reaction for a period of 10–100 s (the uncatalyzed reaction needs around 60–100 s in the assay conditions, whereas the catalyzed ones are of around 6–10 s). The CO_2 concentrations ranged from 1.7 to 17 mM for the determination of the kinetic parameters. The uncatalyzed rates were determined in the same manner and subtracted from the total observed rates. Enzyme concentrations in the assay system were about 16 nM for all the enzymes considered in the present study.

Results and discussion

16S rDNA sequence analysis and characterization of the bacteria

The result of the microbial growth on the B-4 agar plates led to the observation that the B13 bacterial isolate was effectively




Figure 3. Evolutionary relationship analysis, by the neighbor-joining method, of 21 known CAs from prokaryotes, in comparison to *Enterobacter* sp. B13-CA (accession numbers are given in parenthesis). Scale bar represents 0.05 substitutions per nucleotide position.

Figure 4. Multiple amino acid sequence alignment of 21 known β -class CAs: *Enterobacter* sp. B13-CA, *E. coli* str. K-12 substr. MG1655can, *Ralstonia eutropha* can, *Enterobacter* sp. RS1 yadF, *Klebsiella pneumoniae* can, *H. influenzae* can, *C. freundii* yadF, *P. aeruginosa* PAO1 CynT, *P. aeruginosa* PAO1 CA, *S. elongatus* PCC 7942 icfA, *B. suis* 1330 CA, *E. coli* cynT, *M. thermautotrophicus* cab, *S. enterica* mig-5, *P. carotovorum* cah, *H. pylori* CynT, *H. influenzae* Rd KW20 CA, *B. pseudomallei* CA, *P. gingivalis* CD, *V. cholerae* CA, *S. pneumoniae* GA17570 CA and *M. tuberculosis* Rv1284. Triangles indicates the zinc ion-binding residues. Other active side conserved residues were shaded in blue (red letters). The figure was drawn with ESPript.

```

Ent_sp._B13_CA      20      30      40      50      60
E.coli_can          EEDPGFFGKLAQA  N RFLWIG  GS  VP  .AERLTGLEP  EP  VHR
R.eutropha_can     EEDPGFFKLAQA  K RFLWIG  GS  VP  .AERLTGLEP  EL  VHR
Ent_sp._RS1_yadF   EEDPTFFMRLANQ  A EYLWIG  GS  VP  .ANQILGLAP  EV  VHR
K.pneumoniae_can  EEDPGFFKLAQA  N RFLWDG  GS  VP  .AERLTGLEP  EL  VHR
H.influenzae_can  EEDPGFFKLSQT  K RFLWIG  GS  VP  .AERLTGLEP  EL  VHR
C.freundii_yadF    EENSTYFKEADH  T HYLWIG  GS  VP  .AEKLTNLEP  EL  VHR
P.aeruginosa_PAO1_CynT  PARSLQFKSLATR  A KALFIA  GS  VV  .PELLTQREP  EL  VIR
P.aeruginosa_PAO1_CA  QEDPDDFFAKLARQ  T EYLWIG  GS  VP  .ANEIVGMPL  DL  VHR
S.elongatus_PCC_7942_icfA  PSHRDLFEQFAK  H RVLFIT  GS  ID  .PNLITQSGM  EL  VIR
B.suis_1330_CA     EKDPEYFSLRSS  R EFLWIG  GS  VP  .ANVVTGLOP  EV  VHR
E.coli_cynT        PKRRAFLFKQLATQ  S RTLFIS  GS  LV  .PELVITQREP  DL  VIR
M.thermautotrophicus_cab  NQDFRFRDLSDLK  S KLCIIT  M  LD  LLDLLEALRIGR  DA  VIK
S.enterica_mig-5   HDYLAQKRNSIAG  Y AAVILS  I  AP  .AEIVLDAGI  ET  NSR
P.carotovorum_cah  LSPDFSLCETGKN  S INIRQA  N  DP  LQLAFQSGTQQ  IN  GHT
H.pylori_CynT     DYASGFKGFLPMP  S HIATVA  M  LD  .VYRMLGIKE  EA  VIR
H.influenzae_Rd_KW20_CA  EENSTYFKEADH  T HYLWIG  GS  VP  .AEKLTNLEP  EL  VHR
B.pseudomallei_CA  ADDPQYFSLRDLQ  A EYLWIG  GS  VP  .ANQIITGLP  EV  VHR
P.gingivalis_CD   RDLNAQAVAGLEG  F EAILLS  I  VP  .VEYIFDKGI  DL  VGR
V.cholerae_CA     AETPEYFAKLAGK  N DFLWIG  A  VP  .AERLTGLYS  EL  VHR
S.pneumoniae_GA17570_CA  YVALHGQLNLPK  K RVALVT  M  LH  .VAQALGLAL  DA  ILR

```

```

Ent_sp._B13_CA      70      80      90      100
E.coli_can          NVANLVIHTDLN  . . . CLSVVO  AVDV  VEVEHIIICG  YG  GGVQAA
R.eutropha_can     NVANLVIHTDLN  . . . CLSVVO  AVDV  VEVEHIIICG  YG  GGVQAA
Ent_sp._RS1_yadF   NVANLVIHTDLN  . . . CLSVVO  AVDV  VEVEHIIICG  YG  GGVQAA
K.pneumoniae_can  NVANLVIHTDLN  . . . CLSVVO  AVDV  VEVEHIIICG  YG  GGVQAA
H.influenzae_can  NVANQVIHTDFN  . . . CLSVVO  AVDV  KIEHIIICG  TN  GGIHAA
C.freundii_yadF    NVANLVIHTDLN  . . . CLSVVO  AVDV  VEVEHIVCG  YG  GGVQAA
P.aeruginosa_PAO1_CynT  NAGNIVPGYGPQ  G VGSASVE  AVAV  VCGDIVVCG  SD  GAMG  A
P.aeruginosa_PAO1_CA  NVANVVLHTDLN  . . . CLSVVO  AVDV  KVKHILVTG  YG  GGVRAA
S.elongatus_PCC_7942_icfA  NAGNLIPIPPGA  ANG  GEGASIE  AIAA  NIEHVVCG  SH  GAMKGL
B.suis_1330_CA     NVANLVHRRADLN  . . . LLSVLE  AVGV  EIKHIVCG  YG  GGVRAA
E.coli_cynT        NAGNIVPSYGP  EPG  G VGSASVE  AVAA  RVSDIVCG  SN  GAMT  A
M.thermautotrophicus_cab  NAGNIVDDGVIR  . . . SAA  AIA  Y  GVNEIIVG  TD  GMARLD
S.enterica_mig-5   VAGNISNRDMLG  . . . SME  ACAV  GAKVVLVIG  TR  GAVRCA
P.carotovorum_cah  IQVNVSPGNTLL  LDNETFTLQQ  HFHA  SENEIDGKQ  PL  GHFVYK
H.pylori_CynT     NAGCVVTDVIR  . . . SLA  SQRL  GTREIILLH  TD  GMLTFT
H.influenzae_Rd_KW20_CA  NVANQVIHTDFN  . . . CLSVVO  AVDV  KIEHIIICG  TN  GGIHAA
B.pseudomallei_CA  NVANVVLHTDLN  . . . CLSVVO  AVDV  KVKHVMVVG  YG  SGKVNA
P.gingivalis_CD   VAGNVVDDHMLG  . . . SLE  ACEV  GSKVLLVLG  ED  GAKSA
V.cholerae_CA     NVANQVIHTDLN  . . . CLSVVO  AVDV  QVKHIVCG  YG  GGVTA
S.pneumoniae_GA17570_CA  NAGGRVTE MIR  . . . SLV  SQQQ  GTREIVVLH  TD  GAQTFE

```

```

Ent_sp._B13_CA      110     120     130     140
E.coli_can          VENTELG  . . . LIDN  LLHIRDIWFKHSSLGEMP  .QERRDLTLCLE
R.eutropha_can     VENPELG  . . . LINN  LLHIRDIWFKHSSLGEMP  .QERRDLTLCLE
LKRERIG  . . . LADN  LRHVDRVDADKHEAYLGLTLREDDAHTRLCEL
Ent_sp._RS1_yadF   .NPELG  . . . LIDN  LLHIRDIWFKHSSLGEMP  .QERSLDTLCLE
K.pneumoniae_can  VENPELG  . . . LIDN  LLHIRDIWFKHSSLGEMP  .EDRDLTLCLE
H.influenzae_can  MADKDLG  . . . LINN  LLHIRDIWFKHSSLGEMP  .PEKRAMDLTKI
C.freundii_yadF    IENKEQG  . . . LIDN  LLHIRDIWFKHSSLGEMP  .QERRMDTLCLE
P.aeruginosa_PAO1_CynT  IASCACLDQLPA  VAG  LHHAAEAR  AMNSAHEHAS  EAARLDALVRH
P.aeruginosa_PAO1_CA  LHNDQLG  . . . LIDG  WLRISRDLAYEYRHEQLPTEERVDRLCEL
S.elongatus_PCC_7942_icfA  LKLNQLQEDMPL  VYD  LQHAQAQRLRLVLDNYSGYE  TDDLVEILVAE
B.suis_1330_CA     MDGYGHG  . . . IIDN  WLPRIADIAQANQALDTEIENQDRDLRCEL
E.coli_cynT        IASCQCMDHMPA  VSH  LRYADSR  VVNEARPHSD  LPSKAAAMVRE
M.thermautotrophicus_cab  EDLIVSR  . . . . . MRELGVEEVIENFS  IDVLNVPVDEEE
S.enterica_mig-5   IDNAELG  . . . NLTG  LDEIKPAIAKTEYS  GERKGSNYDFVDAVARK
P.carotovorum_cah  NADGALT  . . . VIA  MFQEGEANPQLATAWQQIPARVDQAEDEVRT
H.pylori_CynT     DDDFKR  . . . . . AIQDET  GIRPTWS  PESYDPAE
H.influenzae_Rd_KW20_CA  MADKDLG  . . . LINN  LLHIRDIWFKHSSLGEMP  .PEKRAMDLTKI
B.pseudomallei_CA  LHNRRVG  . . . LADN  LLHVQDVREKHAALLEDWPLGEARYRRLIE
P.gingivalis_CD   IKGVMG  . . . NITS  LMEIKPSVEATQY  MGERTYANKEFADAVVKE
V.cholerae_CA     IDNPFQ  . . . LINN  LLHIRDY  LKHREYLDKMP  AEDRSDKLAIE
S.pneumoniae_GA17570_CA  NEFPQE  . . . . . YLKEEL  GVDVSDQDFL  PFDQTEE

```

```

Ent_sp._B13_CA      150     160     170     180     190
E.coli_can          NVMEQVYNLGH  STIMQSAWKR  GKQVSIHGWA  G  HD  LLRNLEVTAT
R.eutropha_can     NVIEQVYNVCQ  TTVLQDAWSR  GQAVTVHGWW  G  SD  LLRLDGMAS
Ent_sp._RS1_yadF   NVMEQVYNLGH  STIMRSWKR  GKQVSIHGWA  G  HD  LLRNLYVTAT
K.pneumoniae_can  NVMEQVYNLGH  STIMQSAWKR  GKQVSIHGWA  G  HD  LLRLDVTAV
H.influenzae_can  NVAEQVYNLGR  TSIKSAWER  GKLSLHGWW  D  ND  FLVDQGMAT
C.freundii_yadF    NVMEQVYNLGH  STIMRSWKR  GKQVSIHGWA  G  HD  LLRLDVTAT
P.aeruginosa_PAO1_CynT  NVIAQLANL  RTHPCVAR  ALEQGR  LNLHGWW  D  ES  RIDALDGAS
P.aeruginosa_PAO1_CA  NVIIQVANV  SHTSIVQNA  WRHGQSL  SVHGCI  G  KD  LWKNLNVTVS
S.elongatus_PCC_7942_icfA  NVLTQIENL  KTYPIVRS  RLFQGG  LQIFGWI  E  ES  EVLQISRST
B.suis_1330_CA     SVSSQVESL  SRTPVLQ  SAWKDG  KDIIVHGW  M  N  KD  LRDIGDCT
E.coli_cynT        NVIAQLANL  QTHPSV  RLAL  EGR  IALHGWW  D  ES  SIAAFDGT
M.thermautotrophicus_cab  NVIEGVK  RLKSSPLIP  . . . . . ESIGVH  GLI  D  NT  RLKPLYLDED
S.enterica_mig-5   NVELTIEN  IRKNSP  VLKQLE  DEKKIK  IVGSM  H  TG  KVEFFE  V  .
P.carotovorum_cah  VAIQALLP  TSLN  YFRFS  GLST  PPKSE  GIR  L  LD  PVTASAE  Q  .
H.pylori_CynT     DVRSQSL  RRIEV  NPFFVT  . . . . . KHTSL  RGFV  D  AT  KNEVTP  .
H.influenzae_Rd_KW20_CA  NVAEQVYNL  GRSTIK  SAWER  GKLSLHG  WW  D  ND  FLVDQGMAT
B.pseudomallei_CA  NVIEQVYN  VCRTT  IVNDA  WARG  PLTVH  ALV  G  HD  RMRNLG  MAVS
P.gingivalis_CD   NVIIQTM  AVEIR  RDSPI  LKLE  EEG  KIK  IC  GAI  E  ST  KVHFL  .
V.cholerae_CA     NVAEQVYN  LANST  VLQNA  WERG  QAVE  VHF  V  G  ED  RLEYL  GVRCA
S.pneumoniae_GA17570_CA  SVREDM  QLL  IESPLIP  . . . . . DDV  IISG  AI  N  DT  SMTV  VEL  .

```

promoting CaCO_3 precipitation, presumably due to the CA activity present in the bacterium (data not shown). The EzTaxon identification results of 16S rDNA sequence of the B13 strain showed that it possesses a high percentage similarity with bacteria belonging to *Enterobacteriaceae*. Among various such bacteria, strain B13 was 99.04% identical to *Enterobacter xiangfangensis*

strain 10–17, 98.63% to *Enterobacter asburiae* strain JCM6051, 98.56% to *Enterobacter cloacae* strain ATCC 13047, 98.45% to *Enterobacter cowanii* CIP 107300, 98.41% to *Klebsiella michiganensis* W14, 98.23% to *Escherichia hermannii* GTC 347, 97.90% to *Escherichia vulneris* ATCC 33821, to 97.89% *Citrobacter youngae* CECT 5335, to 97.77% *Enterobacter*

Table 1. Kinetic parameters for the CO₂ hydration reaction catalyzed by the human cytosolic isozymes hCA I and II (α -class CAs)⁴⁸ and the bacterial β -CAs: PgiCAB (from *P. gingivalis*)²⁸, HpyCA (from *H. pylori*)³⁵, LpCA1 and LpCA2 (from *L. pneumophila*)²⁴ and B13-CA (from *Enterobacter* sp. B13).

Enzyme	Class	k_{cat} (s ⁻¹)	k_{cat}/K_m (M ⁻¹ × s ⁻¹)	K_I (acetazolamide) (nM)
hCA I	α	2.0×10^5	5.0×10^7	250
hCA II	α	1.4×10^6	1.5×10^8	12
PgiCAB	β	2.8×10^5	1.5×10^7	214
HpyCA	β	7.1×10^5	4.8×10^7	40
LpCA1	β	3.4×10^5	4.7×10^7	76.8
LpCA2	β	8.3×10^5	8.5×10^7	72.1
B13-CA	β	4.8×10^5	5.6×10^7	78.9

All the measurements have been made at 20 °C, pH 8.3 by a stopped flow CO₂ hydrase assay method¹⁹. Acetazolamide (AAZ) inhibition data are also provided.

ludwigii EN-119T, 97.73% to *Enterobacter soli* LF7a, 97.67% to *Cedecea davisae* DSM 4568, 97.63% *Cedecea neteri*, 97.60% to *Erwinia aphidicola* GTC 1688, 97.59% to *Enterobacter aerogenes* KCTC 2190 (Figure 1).

Purification of *Enterobacter* sp. B13-CA

The recombinant B13-CA was purified in concentrations ranging between 0.04 and 2.4 mg/ml from 250 ml cultures. The purified enzyme was subjected to SDS-PAGE analysis (Figure 2) which led to the observation of a distinct single band, with the predicted molecular mass of 24.95 kDa (deduced from the amino acid sequence). Data of the SDS-PAGE (Figure 2, lane 2) showed that the purified B13-CA had indeed a molecular weight (for the monomer) of 25 kDa.

Biochemical characterization of *Enterobacter* sp. B13-CA

According to the zymogram analysis, *Enterobacter* sp. B13 CA did not reveal any esterase activity. Furthermore, no esterase activity was observed by using a spectrophotometric assay with α -naphthyl acetate as substrate (data not shown). This is to be expected, since esterase activity was not reported for any β -class CAs, although many putative activated esters have been investigated as possible substrates²⁰. In fact only the α -^{20,21} and η -CAs²² possess esterase activity with activated esters (such as 4-nitrophenyl- or α -naphthyl acetate) as substrates.

Bioinformatic analysis

Considering the sequence of the *can* gene of *Enterobacter* sp. B13, which contains 663 bp, it encodes a protein of 220 amino acid residues. Based on DELTA-BLAST results of this amino acid sequence, *Enterobacter* sp. B13 *can* shows 33% homology with *E. coli cynT* and 95% with *can* (from the same bacterium, i.e. *E. coli*). Phylogenetic analysis of amino acid resemblance revealed that B13-CA is a member of the β -CA class and has a close relationship with *can* genes identified in other bacteria belonging to *Enterobacteriaceae* (Figure 3). A conserved domain database (CDD) (<http://www.ncbi.nlm.nih.gov/Structure/cdd/wrpsb.cgi>) search of *Enterobacter* sp. B13 *can* sequences showed that the active and ion binding site residues of B13-CA are well conserved, as in all β -CAs investigated in detail, such as among others the enzymes from *H. influenzae*, *Citrobacter freundii*, *Pseudomonas aeruginosa*, *Synechococcus elongatus*, *Brucella suis*, *E. coli*, *Methanothermobacter thermautotrophicus*, *Salmonella enterica*, *Pectobacterium carotovorum*, *Helicobacter*

pylori, *Porphyromonas gingivalis*, *Vibrio cholerae*, *Streptococcus pneumoniae*, *S. mutans* and *Mycobacterium tuberculosis Rv1284*^{23–42} (Figure 4).

Catalytic activity of B13-CA

In fact there are two types of β -CAs in organisms all over the phylogenetic tree, which are defined by their pH–catalytic activity profile as well as active site structural configuration^{41–52}. Type I β -CAs display catalytic activity over a broad pH range (6.5–9.0) with the active site zinc tetrahedrally coordinated by three amino acid residues (two Cys and one His) and a hydroxide/water. In contrast, type II β -CAs are catalytically active only at a pH 8 and higher where they adopt a functional active site configuration like that of type I. In fact, below pH 8 they are conformationally self-inactivated by the addition of a fourth amino acid (an Asp residue) which coordinates to the Zn(II) ion as the fourth ligand, and displacing the zinc bound solvent^{39,52}. At pH > 8, the Asp copordinated to Zn(II) makes a salt pair interaction with a conserved Arg residue, which “opens” the active site in the sense that a water molecule/hydroxide ion takes the place of the Asp residue, generating thus the nucleophile responsible for the catalytic activity of these enzymes^{39,48–52}.

As seen from Figure 4, where an alignment of the amino acid sequence of B13-CA with that of other 20 bacterial β -CAs is presented, the new enzyme described here has all the features of a catalytically effective β -CA: (i) the putative Zn(II) ligands, Cys42, Asp44 (in case B13-CA is a type II β -CA), His 98 and Cys101; (ii) The catalytic dyad involved in activation of the zinc-coordinated water molecule/hydroxide ion for catalysis⁵³, represented by the Asp44–Arg46 residues, which resembles in a way the activation of the water molecule in aspartic proteases⁵³.

As seen from Table 1, in which we compared the catalytic activity of the new enzyme reported here, B13-CA with that of other α - and β -class CAs from various organisms, B13-CA shows a significant activity as catalyst for the hydration of CO₂ with formation of protons and bicarbonate. Indeed, B13-CA possesses the following kinetic parameters at 20 °C and pH of 8.3: k_{cat} of 4.8×10^5 s⁻¹ and k_{cat}/K_m of 5.6×10^7 M⁻¹ × s⁻¹. In fact the catalytic activity of the new enzyme is in the same range as those of other β -CAs, such as those from *P. gingivalis* (PgiCAB)²⁸, *H. pylori* (HpyCA)³⁵, or *Legionella pneumophila* (LpCA1 and LpCA2)²⁴, recently characterized by one of our groups. Furthermore, the enzyme was inhibited by the clinically used^{54–57} sulfonamide CA inhibitor (CAI) acetazolamide (AAZ) with an inhibition constant of 78.9 nM, in the same range as the human (h) hCA I isoform, HpyCA, LpCA1 and LpCA2 (which were inhibited with inhibition constants ranging between 40 and 250 nM). Only the physiologically dominant isoform hCA II was better inhibited by this drug, with a K_I of 12 nM (Table 1).

Conclusions

A recombinant CA (EC 4.2.1.1) from the soil-dwelling bacterium *Enterobacter* sp. B13 was cloned and purified by Co²⁺ affinity chromatography. Bioinformatic analysis showed the new enzyme (denominated here B13-CA) to belong to the β -CA class and to possess 95% homology with the ortholog enzyme from *E. coli* encoded by the *can* gene, whereas its sequence homology with the other such enzyme from *E. coli* (encoded by the *cynT* gene) was of 33%. B13-CA was characterized kinetically as a catalyst for carbon dioxide hydration to bicarbonate and protons. The enzyme showed a significant catalytic activity, with the following kinetic parameters at 20 °C and pH of 8.3: k_{cat} of 4.8×10^5 s⁻¹ and k_{cat}/K_m of 5.6×10^7 M⁻¹ × s⁻¹. This activity was potently inhibited by acetazolamide which showed a K_I of 78.9 nM. Although only this compound was investigated for the moment as

B13-CAI, further studies may reveal new classes of inhibitors/activators of this enzyme^{58–60}, which may show biomedical or environmental applications.

Declaration of interest

The authors do not have any conflicts of interest.

References

- Smith KS, Ferry JG. Prokaryotic carbonic anhydrases. *FEMS Microbiol Rev* 2000;24:335–66.
- Hashimoto M, Kato JI. Indispensability of the *Escherichia coli* carbonic anhydrases YadF and CynT in cell proliferation at a low CO₂ partial pressure. *Biosci Biotechnol Biochem* 2003;67:919–22.
- Rodriguez-Navarro C, Jroundi F, Schiro M, et al. Influence of substrate mineralogy on bacterial mineralization of calcium carbonate: implications for stone conservation. *Appl Environ Microbiol* 2012;78:4017–29.
- Capasso C, De Luca V, Carginale V, et al. Biochemical properties of a novel and highly thermostable bacterial α -carbonic anhydrase from *Sulfurihydrogenibium yellowstonense* YO3AOP1. *J Enzyme Inhib Med Chem* 2012;27:892–7.
- Del Prete S, Vullo D, Fisher GM, et al. Discovery of a new family of carbonic anhydrases in the malaria pathogen *Plasmodium falciparum*—the η -carbonic anhydrases. *Bioorg Med Chem Lett* 2014;24:4389–96.
- Sung YC, Fuchs JA. Characterization of the cyn operon in *Escherichia coli* K12. *J Biol Chem* 1988;263:14769–75.
- Guilloton MB, Korte JJ, Lamblin AF, et al. Carbonic anhydrase in *Escherichia coli*. A product of the cyn operon. *J Biol Chem* 1992;267:3731–4.
- Cronk JD, Endrizzi JA, Cronk MR, et al. Crystal structure of *E. coli* beta-carbonic anhydrase, an enzyme with an unusual pH-dependent activity. *Protein Sci* 2001;10:911–22.
- Merlin C, Masters M, McAteer S, Coulson A. Why is carbonic anhydrase essential to *Escherichia coli*? *J Bacteriol* 2003;185:6415–24.
- Ramanan R, Kannan K, Sivanesan SD, Chakrabarti T. Prevalence and phylogenetic relationship of two β -carbonic anhydrases in affiliates of *Enterobacteriaceae*. *Ann Microbiol* 2013;63:1275–82.
- Chirică LC, Elleby B, Jonsson BH, Lindskog S. The complete sequence, expression in *Escherichia coli*, purification and some properties of carbonic anhydrase from *Neisseria gonorrhoeae*. *Eur J Biochem* 1997;244:755–60.
- Joseph S, David WR. *Molecular cloning: a laboratory manual*. Cold Spring Harbor (NY): Cold Spring Harbor Laboratory Press; 2001:2–25.
- Kotwica J, Ciuk MA, Joachimiak E, et al. Carbonic anhydrase activity in the vas deferens of the cotton leafworm—*Spodoptera littoralis* (Lepidoptera: Noctuidae) controlled by circadian clock. *J Physiol Pharmacol* 2006;57:107–23.
- Eminoğlu A, Ülker S, Sandallı C. Cloning, purification and characterization of acetyl xylane esterase from *Anoxybacillus flavithermus* DSM 2641(T) with activity on low molecular-weight acetates. *Protein J* 2015;34:237–42.
- Kim OS, Cho YJ, Lee K, et al. Introducing EzTaxon-e: a prokaryotic 16S rRNA gene sequence database with phylotypes that represent uncultured species. *Int J Syst Evol Microbiol* 2012;62:716–21.
- Bjellqvist B, Hughes GJ, Pasquali C, et al. The focusing positions of polypeptides in immobilized pH gradients can be predicted from their amino acid sequences. *Electrophoresis* 1993;14:1023–31.
- Thompson JD, Higgins DG, Gibson TJ. CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res* 1994;22:4673–80.
- Tamura K, Stecher G, Peterson D, et al. MEGA6: molecular evolutionary genetics analysis version 6.0. *Mol Biol Evol* 2013;30:2725–9.
- Khalifah RG. The carbon dioxide hydration activity of carbonic anhydrase. I. Stop-flow kinetic studies on the native human isoenzymes B and C. *J Biol Chem* 1971;246:2561–73.
- Innocenti A, Scozzafava A, Parkkila S, et al. Investigations of the esterase, phosphatase and sulfatase activities of the cytosolic mammalian carbonic anhydrase isoforms I, II and XIII with 4-nitrophenyl esters as substrates. *Bioorg Med Chem Lett* 2008;18:2267–71.
- Del Prete S, Vullo D, De Luca V, et al. Biochemical characterization of the δ -carbonic anhydrase from the marine diatom *Thalassiosira weissflogii*, TweCA. *J Enzyme Inhib Med Chem* 2014;29:906–11.
- Krungkrai J, Krungrai SR, Supuran CT. Carbonic anhydrase inhibitors: inhibition of *Plasmodium falciparum* carbonic anhydrase with aromatic/heterocyclic sulfonamides—in vitro and in vivo studies. *Bioorg Med Chem Lett* 2008;18:5466–71.
- Vullo D, Del Prete S, Osman SM, et al. Sulfonamide inhibition studies of the gamma-carbonic anhydrase from the oral pathogen *Porphyromonas gingivalis*. *Bioorg Med Chem Lett* 2014;24:240–4.
- Nishimori I, Vullo D, Minakuchi T, et al. Anion inhibition studies of two new β -carbonic anhydrases from the bacterial pathogen *Legionella pneumophila*. *Bioorg Med Chem Lett* 2014;24:1127–32.
- Del Prete S, De Luca V, Scozzafava A, et al. Biochemical properties of a new α -carbonic anhydrase from the human pathogenic bacterium, *Vibrio cholerae*. *J Enzyme Inhib Med Chem* 2014;29:23–7.
- Del Prete S, De Luca V, Vullo D, et al. Biochemical characterization of the γ -carbonic anhydrase from the oral pathogen *Porphyromonas gingivalis*, PgiCA. *J Enzyme Inhib Med Chem* 2014;29:532–7.
- Del Prete S, Isik S, Vullo D, et al. DNA cloning, characterization, and inhibition studies of an alpha-carbonic anhydrase from the pathogenic bacterium *Vibrio cholerae*. *J Med Chem* 2012;55:10742–8.
- Del Prete S, Vullo D, De Luca V, et al. Biochemical characterization of recombinant beta-carbonic anhydrase (PgiCAB) identified in the genome of the oral pathogenic bacterium *Porphyromonas gingivalis*. *J Enzyme Inhib Med Chem* 2015;30:366–70.
- Supuran CT. Bacterial carbonic anhydrases as drug targets: toward novel antibiotics? *Front Pharmacol* 2011;2:34.
- Vullo D, Nishimori I, Minakuchi T, et al. Inhibition studies with anions and small molecules of two novel beta-carbonic anhydrases from the bacterial pathogen *Salmonella enterica* serovar *Typhimurium*. *Bioorg Med Chem Lett* 2011;21:3591–5.
- Joseph P, Ouahrani-Bettache S, Montero JL, et al. A new beta-carbonic anhydrase from *Brucella suis*, its cloning, characterization, and inhibition with sulfonamides and sulfamates, leading to impaired pathogen growth. *Bioorg Med Chem* 2011;19:1172–8.
- Joseph P, Turtaut F, Ouahrani-Bettache S, et al. Cloning, characterization, and inhibition studies of a beta-carbonic anhydrase from *Brucella suis*. *J Med Chem* 2010;53:2277–85.
- Nishimori I, Minakuchi T, Morimoto K, et al. Carbonic anhydrase inhibitors: DNA cloning and inhibition studies of the alpha-carbonic anhydrase from *Helicobacter pylori*, a new target for developing sulfonamide and sulfamate gastric drugs. *J Med Chem* 2006;49:2117–26.
- Nishimori I, Onishi S, Takeuchi H, et al. The alpha and beta classes carbonic anhydrases from *Helicobacter pylori* as novel drug targets. *Curr Pharm Des* 2008;14:622–30.
- Nishimori I, Minakuchi T, Kohsaki T, et al. Carbonic anhydrase inhibitors: the beta-carbonic anhydrase from *Helicobacter pylori* is a new target for sulfonamide and sulfamate inhibitors. *Bioorg Med Chem Lett* 2007;17:3585–94.
- Burghout P, Vullo D, Scozzafava A, et al. Inhibition of the β -carbonic anhydrase from *Streptococcus pneumoniae* by inorganic anions and small molecules: toward innovative drug design of anti-infectives? *Bioorg Med Chem* 2011;19:243–8.
- Guzel O, Maresca A, Scozzafava A, et al. Discovery of low nanomolar and subnanomolar inhibitors of the mycobacterial beta-carbonic anhydrases Rv1284 and Rv3273. *J Med Chem* 2009;52:4063–7.
- Nishimori I, Minakuchi T, Maresca A, et al. The β -carbonic anhydrases from *Mycobacterium tuberculosis* as drug targets. *Curr Pharm Des* 2010;16:3300–9.
- Suarez Covarrubias A, Larsson AM, Hogbom M, et al. Structure and function of carbonic anhydrases from *Mycobacterium tuberculosis*. *J Biol Chem* 2005;280:18782–9.
- Carta F, Maresca A, Covarrubias AS, et al. Carbonic anhydrase inhibitors. Characterization and inhibition studies of the most active beta-carbonic anhydrase from *Mycobacterium tuberculosis*, Rv3588c. *Bioorg Med Chem Lett* 2009;19:6649–54.
- Minakuchi T, Nishimori I, Vullo D, et al. Molecular cloning, characterization, and inhibition studies of the Rv1284 beta-carbonic

- anhydrase from *Mycobacterium tuberculosis* with sulfonamides and a sulfamate. *J Med Chem* 2009;52:2226–32.
42. Capasso C, Supuran CT. Sulfa and trimethoprim-like drugs—antimetabolites acting as carbonic anhydrase, dihydropteroate synthase and dihydrofolate reductase inhibitors. *J Enzyme Inhib Med Chem* 2014;29:379–87.
 43. Migliardini F, De Luca V, Carginale V, et al. Biomimetic CO₂ capture using a highly thermostable bacterial α -carbonic anhydrase immobilized on a polyurethane foam. *J Enzyme Inhib Med Chem* 2014;29:146–50.
 44. Supuran CT, Capasso C. The η -class carbonic anhydrases as drug targets for antimalarial agents. *Expert Opin Ther Targets* 2015;19:551–63.
 45. Capasso C, Supuran CT. An overview of the alpha-, beta- and gamma-carbonic anhydrases from Bacteria: can bacterial carbonic anhydrases shed new light on evolution of bacteria? *J Enzyme Inhib Med Chem* 2015;30:325–32.
 46. Supuran CT. Carbonic anhydrase inhibitors: an editorial. *Expert Opin Ther Pat* 2013;23:677–79.
 47. Capasso C, Supuran CT. Anti-infective carbonic anhydrase inhibitors: a patent and literature review. *Expert Opin Ther Pat* 2013;23:693–704.
 48. Supuran CT. Carbonic anhydrases: novel therapeutic applications for inhibitors and activators. *Nat Rev Drug Discov* 2008;7:168–81.
 49. De Luca V, Del Prete S, Supuran CT, Capasso C. Protonography, a new technique for the analysis of carbonic anhydrase activity. *J Enzyme Inhib Med Chem* 2015;30:277–82.
 50. Supuran CT. Structure-based drug discovery of carbonic anhydrase inhibitors. *J Enzyme Inhib Med Chem* 2012;27:759–72.
 51. Supuran CT. Carbonic anhydrases: from biomedical applications of the inhibitors and activators to biotechnological use for CO(2) capture. *J Enzyme Inhib Med Chem* 2013;28:229–30.
 52. Pinard MA, Lotlikar SR, Boone CD, et al. Structure and inhibition studies of a type II beta-carbonic anhydrase psCA3 from *Pseudomonas aeruginosa*. *Bioorg Med Chem* 2015; 23:4831–8.
 53. Mastrolorenzo A, Rusconi S, Scozzafava A, et al. Inhibitors of HIV-1 protease: current state of the art 10 years after their introduction. From antiretroviral drugs to antifungal, antibacterial and antitumor agents based on aspartic protease inhibitors. *Curr Med Chem* 2007; 14:2734–48.
 54. Supuran CT. Carbonic anhydrase inhibition/activation: trip of a scientist around the world in the search of novel chemotypes and drug targets. *Curr Pharm Des* 2010;16:3233–45.
 55. Supuran CT. Carbonic anhydrase inhibitors. *Bioorg Med Chem Lett* 2010;20:3467–74.
 56. De Simone G, Alterio V, Supuran CT. Exploiting the hydrophobic and hydrophilic binding sites for designing carbonic anhydrase inhibitors. *Expert Opin Drug Discov* 2013;8:793–810.
 57. Alterio V, Di Fiore A, D'Ambrosio K, et al. Multiple binding modes of inhibitors to carbonic anhydrases: how to design specific drugs targeting 15 different isoforms? *Chem Rev* 2012; 112:4421–68.
 58. Maresca A, Carta F, Vullo D, et al. Dithiocarbamates strongly inhibit the β -class carbonic anhydrases from *Mycobacterium tuberculosis*. *J Enzyme Inhib Med Chem* 2013;28:407–11.
 59. Scozzafava A, Passaponti M, Supuran CT, Gulcin I. Carbonic anhydrase inhibitors: guaiacol and catechol derivatives effectively inhibit certain human carbonic anhydrase isoenzymes (hCA I, II, IX and XII). *J Enzyme Inhib Med Chem* 2015;30:586–91.
 60. Alp C, Maresca A, Alp NA, et al. Secondary/tertiary benzenesulfonamides with inhibitory action against the cytosolic human carbonic anhydrase isoforms I and II. *J Enzyme Inhib Med Chem* 2013;28: 294–8.